

ON THE $|N, p_n, q_n|_k$ SUMMABILITY FACTORS OF INFINITE SERIES

Debadutta Mohanty

Abstract

In this paper a theorem on generalized Nörlund Summability Factors has been proved which generalizes some earlier factor theorems on

$|N, p_n|_k$ ([5]), $|C, 1|_k$ ([4]) and $|\bar{N}, p_n|_k$ ([1]) for $k > 1$.

1. INTRODUCTION

Given any series $\sum a_n$, if there exists a sequence $\{\lambda_n\}$ such that $\sum_{n=0}^{\infty} a_n \lambda_n$ is summable by a method A, then we say that $\{\lambda_n\}$ is a summability factor for the method A. Results establishing theorems on summability factors are called factor theorems.

In this section we introduce some notations, conventions and definitions which are to be used in this paper. Let $\sum_0^{\infty} a_n$ be an infinite series with sequences of partial sums $\{s_n\}$.

Let $p = \{p_n\}$ be a positive non-increasing sequence of real numbers such that

$$p_n = \sum_{i=0}^n p_i \rightarrow \infty \text{ as } n \rightarrow \infty; \tag{1.1}$$

and

$$p_{-i} = p_{-i} = 0 \text{ for } i \geq 1.$$

For a positive real sequence $q = (q_n)$, we define an increasing sequence (r_n) by

Mathematics Subject Classification: 40D15, 40F 05, 40G99

Key Words: Mathematics Subject Classification: 40D15, 10F05, 40G99.

$$r_n = (p * q)_n = \sum_{i=0}^n p_{n-i} q_{i \rightarrow \infty}, \text{ as } n \rightarrow \infty \quad (1.2)$$

where $q_n = 0(1)$ and $1 = 0(q_n)$ as $n \rightarrow \infty$ and $q_{-i} = Q_{-i} = r_{-i} = 0$ for $i \geq 1$, $*$ denotes the convolution product.

The (N, p_n, q_n) -transform of the sequence (s_n) is defined by ([2]).

$$t_n = \frac{1}{r_n} \sum_{i=0}^n p_{n-i} q_i s_i. \quad (1.3)$$

The series $\sum a_n$ is said to be $|N, p_n, q_n|_k$ summable for $k \geq 1$, if

$$\sum_{n=1}^{\infty} \left(\frac{r_n}{q_n} \right)^{k-1} |t_n - t_{n-1}|^k < \infty. \quad (1.4)$$

If (d_n) be the sequence of $(C, 1)$ -transform of the sequence (n, a_n) , then

$$d_n = \frac{1}{n+1} \sum_{k=0}^n k a_k = \frac{1}{n+1} \sum_{k=1}^n k a_k \text{ as } a_0 = 0. \quad (1.5)$$

2. NOTATIONS

We use the following notations

$$A_j^k = \sum_{j=i}^k p_{k-j} q_j = p_{k-i} q_i + p_{k-i-1} q_{i+1} + \dots + p_0 q_k$$

so that

$$\begin{aligned} A_j^k &= \sum_{j=i}^k p_{k-j} q_j, \quad 0 \leq i < k \\ &= A_i^n, \quad k \geq n \\ &= 0, \quad i > k \end{aligned} \quad (2.1)$$

and

$$B_i^k = \sum_{j=0}^k p_{n-j} q_{j-i} = p_{n-i} q_0 + p_{n-i-1} q_1 + \dots + p_{n-k} q_{k-i}$$

so that

$$\begin{aligned} B_i^k &= \sum_{j=i}^k p_{n-j} q_{j-i}, \quad 0 \leq i < k < n \\ &= B_i^n, \quad k \geq n \\ &= 0, \quad i > k \end{aligned} \tag{2.2}$$

we find the relations as

$$B_k^n = r_{n-k} = A_0^{n-k}$$

and

$$B_i^k + A_{k+1-i}^{n-i} = (p * q)_{n-i} = r_{n-i}.$$

In particular, if $i = 0$, then

$$B_0^k + A_{k+1}^n = r_n.$$

3. The object of this note is to prove the following theorem on $|N, p_n, q_n|_k$ summability.

Theorem: Let (p_n) , (q_n) and (r_n) be the sequences satisfying (1.1) and (1.2). If (X_n) is a positive monotonic non-decreasing sequence and (λ_n) be any sequence such that

$$\lambda_m X_m = o(1) \text{ as } m \rightarrow \infty \tag{3.1}$$

$$\sum_{n=1}^m n X_n |\Delta^2 \lambda_n| = o(1), \quad m \rightarrow \infty \tag{3.2}$$

$$\sum_{n=1}^m \frac{q_n}{r_n} |d_n|^k = o(X_m), \quad m \rightarrow \infty \tag{3.3}$$

and

$$r_n = o(n, q_n) \text{ as } n \rightarrow \infty \tag{3.4}$$

where d_n is the n -th $(C, 1)$ transform of the sequence (na_n) , then the series $\sum a_n \lambda_n$ is summable $|N, p_n, q_n|_k$, $k \geq 1$.

It may be noticed that under the conditions of the theorem, we have that

$$\Delta \lambda_n = \lambda_n - \lambda_{n+1} \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (3.5)$$

4. We require the following Lemmas to prove the theorem.

It is easy to prove the following two equalities as

$$r_{n-1} A_i^n - r_n A_i^{n-1} = B_i^i A_i^n - B_0^{i-1} A_i^{n-1} \quad (4.1)$$

and

$$\begin{aligned} & \left| B_1^i A_i^n + B_0^i A_{i+1}^{n-1} - B_0^{i-1} A_i^{n-1} - B_1^{i+1} A_{i+1}^n \right| \quad (4.2) \\ &= \left| q_i (r_{n-1} p_{n-i} - r_n p_{n-i-1}) \right| \\ &= q_i (r_n p_{n-i-1} - r_{n-1} p_{n-i}). \end{aligned}$$

Lemma 1: Under the relations given above, we find

$$\frac{1}{q_n r_{n-1}} \sum_{i=1}^n \left| B_i^i A_i^n + B_0^i A_{i+1}^{n-1} - B_0^{i-1} A_i^{n-1} - B_1^{i+1} A_{i+1}^n \right| = 0(1), \quad (4.3)$$

as $n \rightarrow \infty$.

Proof: From (4.2)

$$\begin{aligned} \text{L.H.S} &= \frac{1}{q_n r_{n-1}} \sum_{i=1}^{n-1} q_i (r_n p_{n-i-1} - r_{n-1} p_{n-i}) \quad (4.4) \\ &= \frac{r_n}{q_n r_{n-1}} \sum_{i=1}^{n-1} q_i p_{n-i-1} - \frac{1}{q_n} \sum_{i=1}^{n-1} q_i p_{n-i} \\ &= \frac{r_n}{q_n r_{n-1}} (r_{n-1} - p_{n-1} q_0) - \frac{1}{q_n} (r_n - p_n q_0 - p_0 q_n) \\ &= \frac{p_n q_0}{q_n} + p_0 - \frac{r_n p_{n-1} q_0}{q_n r_{n-1}} \\ &= \frac{q_0}{q_n} \left(p_n - \frac{r_n}{r_{n-1}} p_{n-1} \right) + p_0 \end{aligned}$$

$$\begin{aligned} &\leq \frac{q_0}{q_n} (p_n - p_{n-1}) + p_0 = o(1) \text{ as } n \rightarrow \infty \\ &= \text{R.H.S.} \end{aligned}$$

Hence the Lemma.

We assume here that

$$\frac{1}{q_n r_{n-1}} \sum_{i=1}^{n-1} \frac{q_i (B_1^i A_i^n - B_0^{i-1} A_{i+1}^{n-1})}{r_i} = o(1) \tag{4.5}$$

and

$$\frac{1}{q_n r_{n-1}} \sum_{i=1}^{n-1} \frac{q_i (B_1^i A_{i+1}^n - B_0^i A_{i+1}^{n-1})}{r_i} = o(1) \text{ as } n \rightarrow \infty. \tag{4.6}$$

Lemma 2: ([1], Lemma)

Suppose (X_n) is a positive non-decreasing sequence such that (3.1) and (3.2) hold. Then

$$\sum_{n=1}^{\infty} X_n |\Delta \lambda_n| < \infty \tag{4.7}$$

and

$$n X_n |\Delta \lambda_n| = o(1), \text{ as } n \rightarrow \infty.$$

Lemma 3: As the notations defined above, we get

$$\sum_{n=i+1}^{m+1} \frac{B_1^i A_i^n - B_0^{i-1} A_i^{n-1}}{r_n r_{n-1}} = o(1), \text{ as } m \rightarrow \infty.$$

Proof: We have

$$\begin{aligned} \sum_{n=i+1}^{m+1} \frac{B_1^i A_i^n - B_0^{i-1} A_i^{n-1}}{r_n r_{n-1}} &= \sum_{n=i+1}^{m+1} \frac{r_n B_1^i - r_{n-1} B_0^{i-1}}{r_n r_{n-1}} \\ &= \Lambda(m, i), \text{ say.} \end{aligned}$$

Clearly $\Lambda(m, i)$ is one positive and decreasing sequence for $1 \leq i \leq m$, hence,

$$\sup_{1 \leq i \leq m} \Lambda(m, i) = \Lambda(m, 1)$$

$$\begin{aligned}
&= \sum_{n=2}^{m+1} \frac{B_1^1 r_n - B_0^0 r_{n-1}}{r_n r_{n-1}} = \sum_{n=2}^{m+1} \left(\frac{B_1^1}{r_{n-1}} - \frac{B_0^0}{r_n} \right) \\
&= \sum_{n=2}^{m+1} \left(\frac{p_{n-1} q_0}{r_{n-1}} - \frac{p_n q_0}{r_n} \right) = \left(\frac{p_1}{r_1} - \frac{p_{m+1}}{r_{m+1}} \right) q_0 \rightarrow \frac{p_1 q_0}{r_1} \text{ as } m \rightarrow \infty.
\end{aligned}$$

Thus $\Lambda(m, i) = 0(1)$ as $m \rightarrow \infty$ for each $i \leq m$; this completes the proof of Lemma.

Similarly we can prove for each i ,

$$\sum_{n=i+1}^{m+1} \frac{B_1^i A_{i+1}^n - B_0^i A_{i+1}^{n-1}}{r_n r_{n-1}} = 0(1), \text{ as } m \rightarrow \infty. \quad (4.9)$$

5. PROOF OF THE THEOREM

Let T_n be the n -th (N, p_n, q_n) transform of the series $\sum a_n \lambda_n$, then

$$\begin{aligned}
T_n &= \frac{1}{r_n} \sum_{i=0}^n p_{n-i} q_i \sum_{k=0}^i a_k \lambda_k \\
&= \frac{1}{r_n} \sum_{k=0}^n a_k \lambda_k \sum_{i=k}^n p_{n-i} q_i = \frac{1}{r_n} \sum_{k=0}^n a_k \lambda_k A_k^n.
\end{aligned}$$

Then for $n \geq 1$, we get, by Abel's Transformation

$$\begin{aligned}
T_n - T_{n-1} &= \frac{1}{r_n} \sum_{i=1}^n A_i^n a_i \lambda_i - \frac{1}{r_{n-1}} \sum_{i=1}^{n-1} A_i^{n-1} a_i \lambda_i \\
&= \sum_{i=1}^n \left(\frac{A_i^n}{r_n} - \frac{A_i^{n-1}}{r_{n-1}} \right) a_i \lambda_i, \text{ since } A_n^{n-1} = 0 \\
&= \frac{1}{r_n r_{n-1}} \sum_{i=1}^n (r_{n-1} A_i^n - A_i^{n-1} r_n) a_i \lambda_i \\
&= \frac{1}{r_n r_{n-1}} \sum_{i=1}^n (B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) a_i \lambda_i \text{ (by (4.1))}
\end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{r_n r_{n-1}} \sum_{i=1}^n \frac{(B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \lambda_i}{i} i a_i \\
 &= \frac{1}{r_n r_{n-1}} \left\{ \sum_{i=1}^{n-1} \Delta \frac{(B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \lambda_i}{i} \sum_{k=1}^i k a_k \right. \\
 &\quad \left. + \frac{(B_1^n A_n^n - B_0^{n-1} A_n^{n-1}) \lambda_n}{n} \sum_{k=1}^n k \cdot a_k + 0 \right\} \\
 &= \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \left[\frac{(B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \lambda_i}{i} - \frac{(B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}) \lambda_{i+1}}{i+1} \right] (i+1) d_i \\
 &\quad + \frac{1}{r_n r_{n-1}} \frac{(B_1^n A_n^n - B_0^{n-1} \cdot 0)}{n} (n+1) d_n.
 \end{aligned}$$

As $\Delta \lambda_i = \lambda_i - \lambda_{i+1}$, we have, by using (4.2)

$$\begin{aligned}
 T_n - T_{n-1} &= \frac{(n+1) p_0 q_n \lambda_n d_n}{n r_n} + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \left\{ \frac{(B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \lambda_i}{i} \right. \\
 &\quad \left. - \frac{B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}}{(i+1)} (\lambda_i - \Delta \lambda_i) \right\} (i+1) d_i \\
 &= \frac{(n+1) p_0 q_n \lambda_n d_n}{n r_n} + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \left\{ (B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \right. \\
 &\quad \left. - (B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}) \right\} \lambda_i d_i + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \frac{1}{i} (B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \lambda_i d_i \\
 &\quad + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} (B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}) \Delta \lambda_i d_i \\
 &= \frac{(n+1) p_0 q_n \lambda_n d_n}{n r_n} + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} q_i (r_{n-i} p_{n-i} - r_n - p_{n-i-1}) \lambda_i d_i
 \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \frac{1}{i} \left(B_1^i A_i^n - B_0^{i-1} A_i^{n-1} \right) \lambda_i d_i + \\
& + \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \left(B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1} \right) \Delta \lambda_i d_i \\
& = T_{n1} + T_{n2} + T_{n3} + T_{n4}, \text{ say.}
\end{aligned}$$

To prove the Theorem, it is sufficient to show, by Minkowski's inequality, that

$$\sum_{n=1}^{\infty} \left(\frac{r_n}{q_n} \right)^{k-1} |T_{nj}|^k < \infty \text{ for } j = 1, 2, 3, 4.$$

Now, by using Abel's transformation

$$\begin{aligned}
I_1 & = \sum_{n=1}^{m+1} \left(\frac{r_n}{q_n} \right)^{k-1} |T_{n1}|^k \\
& = \sum_{n=1}^{m+1} \frac{p_0^k q_n |\lambda_n|^{k-1} |\lambda_n| |d_n|^k}{r_n} \left(\frac{n+1}{n} \right)^k \\
& = O(1) \sum_{n=1}^{m+1} \frac{q_n |\lambda_n| |d_n|^k}{r_n} \\
& = O(1) \left[\sum_{n=1}^m \Delta |\lambda_n| \sum_{i=1}^n \frac{q_i |d_i|^k}{r_i} + |\lambda_m| \sum_{i=1}^m \frac{q_i |d_i|^k}{r_i} \right] \\
& = O(1) \sum_{n=1}^m \Delta |\lambda_n| X_n + O(1) |\lambda_m| X_m \\
& = O(1) \sum_{n=1}^m \Delta |\lambda_n| X_n + O(1) \\
& = O(1) \text{ as } m \rightarrow \infty
\end{aligned}$$

by virtue of Lemma 2 and hypothesis.

Next, by using Holder's inequality and (4.4), we get

$$\begin{aligned}
 I_2 &= \sum_{n=1}^{m+1} \left(\frac{r_n}{q_n} \right)^{k-1} |T_{n2}|^k \\
 &\leq \sum_{n=2}^{m+1} \frac{1}{q_n^{k-1} r_n r_{n-1}} \left\{ \sum_{i=1}^{n-1} q_i (r_n p_{n-i-1} - r_{n-1} p_{n-i}) |\lambda_i| |d_i| \right\}^k \\
 &\leq \sum_{n=2}^{m+1} \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} q_i (r_n p_{n-i-1} - r_{n-1} p_{n-i}) |\lambda_i|^k |d_i|^k \\
 &\quad \times \left\{ \frac{1}{q_n r_n} \sum_{i=1}^{n-1} q_i (r_n p_{n-i-1} - r_{n-1} p_{n-i}) \right\}^{k-1} \\
 &= O(1) \sum_{i=1}^m q_i |\lambda_i|^k |d_i|^k \sum_{n=i+1}^{m+1} \frac{r_n p_{n-i-1} - r_{n-1} p_{n-i}}{r_n r_{n-1}} \\
 &= O(1) \sum_{i=1}^m q_i |\lambda_i|^{k-1} |\lambda_i| |d_i|^k \sum_{n=i+1}^{m+1} \left(\frac{p_{n-i-1}}{r_{n-1}} - \frac{p_{n-i}}{r_n} \right) \\
 &= O(1) \sum_{i=1}^m q_i |\lambda_i| |d_i|^k \left(\frac{p_0}{r_i} - \frac{p_{m+1-i}}{r_{m+1}} \right) \\
 &= O(1) \sum_{i=1}^m \frac{|\lambda_i| q_i |d_i|^k}{r_i} + O(1) \sum_{i=1}^m \frac{|\lambda_i| q_i |d_i|^k}{r_{m+1}} \\
 &= O(1) \text{ as } m \rightarrow \infty, \text{ from } I_1.
 \end{aligned}$$

Also, from the fact that $r_n = O(nq_n)$, we find

$$\begin{aligned}
 I_3 &= \sum_{n=1}^{m+1} \left(\frac{r_n}{q_n} \right)^{k-1} |T_{n3}|^k \\
 &= \sum_{n=2}^{m+1} \left(\frac{r_n}{q_n} \right)^{k-1} \left| \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \frac{1}{i} (B_1^i A_i^n - B_0^{i-1} A_i^{n-1}) \lambda_i d_i \right|^k
 \end{aligned}$$

$$\begin{aligned}
&= 0(1) \sum_{n=2}^{m+1} \frac{1}{q_n^{k-1} r_n r_{n-1}} \left\{ \sum_{i=1}^{n-1} \frac{q_i (B_1^i A_i^n - B_0^{i-1} A_i^{n-1})}{r_i} |\lambda_i| |d_i| \right\}^k \\
&= 0(1) \sum_{n=2}^{m+1} \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \frac{q_i (B_1^i A_i^n - B_0^{i-1} A_i^{n-1})}{r_i} |\lambda_i|^k |d_i|^k \\
&\quad \times \left\{ \frac{1}{q_n r_{n-1}} \sum_{i=1}^{n-1} \frac{q_i (B_1^i A_i^n - B_0^{i-1} A_i^{n-1})^{k-1}}{r_i} \right\}
\end{aligned}$$

This is due to Hölder's inequality. Now from (4.5) and Lemma 3, we get

$$\begin{aligned}
I_3 &= 0(1) \sum_{n=2}^{m+1} \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} \frac{q_i (B_1^i A_i^n - B_0^{i-1} A_i^{n-1})}{r_i} |\lambda_i|^k |d_i|^k \\
&= 0(1) \sum_{n=1}^m \frac{q_i |\lambda_i|^k |d_i|^k}{r_i} \sum_{n=i+1}^{m+1} \frac{(B_1^i A_i^n - B_0^{i-1} A_i^{n-1})}{r_n r_{n-1}} \\
&= 0(1) \sum_{i=1}^m \frac{q_i |\lambda_i| |\lambda_i|^{k-1} |d_i|^k}{r_i} \\
&= 0(1) \sum_{i=1}^m \frac{|\lambda_i| q_i |d_i|^k}{r_i} \\
&= 0(1) \text{ as } m \rightarrow \infty, \text{ from } I_1.
\end{aligned}$$

Lastly, from (4.6) and (4.9) and by applying Hölder's inequality, we have that

$$\begin{aligned}
I_4 &= \sum_{n=1}^{m+1} \left(\frac{r_n}{q_n} \right)^{k-1} |T_{n4}|^k \\
&= \sum_{n=2}^{m+1} \left(\frac{r_n}{q_n} \right)^{k-1} \left| \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} (B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}) \Delta \lambda_i d_i \right|^k \\
&\leq \sum_{n=2}^{m+1} \frac{1}{q_n^{k-1} r_n r_{n-1}} \left\{ \sum_{i=1}^{n-1} \frac{i (B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1})}{i} |\Delta \lambda_i| |d_i| \right\}^k
\end{aligned}$$

$$\begin{aligned}
 &= 0(1) \sum_{n=2}^{m+1} \frac{1}{q_n^{k-1} r_n r_{n-1}^k} \left\{ \sum_{i=1}^{n-1} \frac{i(B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}) q_i}{r_i} |\Delta \lambda_i| |d_i| \right\}^k \\
 &= 0(1) \sum_{n=2}^{m+1} \frac{1}{r_n r_{n-1}} \sum_{i=1}^{n-1} i^k \frac{(B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1})}{r_i} |\Delta \lambda_i|^k |d_i|^k \\
 &\quad \times \left\{ \frac{1}{q_n r_{n-1}} \sum_{i=1}^{n-1} \frac{B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1}}{r_i} \right\}^{k-1} \\
 &= 0(1) \sum_{i=1}^m \frac{q_i}{r_i} i^k |\Delta \lambda_i|^k |d_i|^k \sum_{n=i+1}^{m+1} \frac{(B_1^{i+1} A_{i+1}^n - B_0^i A_{i+1}^{n-1})}{r_n r_{n-1}} \\
 &= 0(1) \sum_{i=1}^m (i|\Delta \lambda_i|)^{k-1} (i|\Delta \lambda_i|) \frac{q_i}{r_i} |d_i|^k \\
 &= 0(1) \sum_{i=1}^m i|\Delta \lambda_i| \frac{q_i |d_i|^k}{r_i} \\
 &= 0(1) \left[\sum_{i=1}^{m-1} \Delta(i|\Delta \lambda_i|) \sum_{v=1}^i \frac{q_v |d_v|^k}{r_v} + m|\Delta \lambda_m| \sum_{i=1}^m \frac{q_i |d_i|^k}{r_i} \right] \\
 &= 0(1) \sum_{i=1}^{m-1} \Delta(i|\Delta \lambda_i|) X_i + 0(1) m|\Delta \lambda_m| X_m \\
 &= 0(1) \sum_{i=1}^{m-1} i X_i |\Delta^2 \lambda_i| + 0(1) \sum_{i=1}^{m-1} |\Delta \lambda_{i+1}| X_i + 0(1) \\
 &= 0(1), \text{ as } m \rightarrow \infty
 \end{aligned}$$

by virtue of the hypothesis and Lemma 2.

This completes the proof of the theorem.

Corollary 1 ([1]): Let (q_n) be a sequence of positive number such that

$$Q_n = 0(n, q_n) \text{ as } n \rightarrow \infty.$$

If (x_n) is a positive monotonic non decreasing sequence such that

$$\lambda_m X_m = o(1) \text{ as } m \rightarrow \infty,$$

$$\sum_{n=1}^{\infty} n X_n |\Delta^2 \lambda_n| = o(1)$$

and

$$\sum_{n=1}^{\infty} \frac{q_n}{Q_n} |t_n|^k = o(X_m) \text{ as } m \rightarrow \infty,$$

where t_n is the n -th $(C, 1)$ transform of (na_n) , then the series $\sum a_n \lambda_n$ is summable $|\bar{N}, q_n|_k$, $k \geq 1$.

Taking $p_n = 1$ for all n in the theorem we get the corollary. In addition to this if we take $q_n = 1$ for all values of n we find,

Corollary 2([4]): If (X_n) is a positive monotonic non-decreasing sequence such that

$$\lambda_m X_m = o(1) \text{ as } m \rightarrow \infty$$

$$\sum_{n=1}^m n X_n |\Delta^2 \lambda_n| = o(1)$$

and

$$\sum_{n=1}^m \frac{1}{n} |t_n|^k = o(X_m) \text{ as } m \rightarrow \infty,$$

then the series $\sum a_n \lambda_n$ is summable $|C, 1|_k$, $k \geq 1$.

Corollary 3 ([5]): Let (p_n) be a monotonic decreasing sequence such that

$$P_n = o(n) \text{ as } n \rightarrow \infty.$$

If (X_n) is a positive monotonic non-decreasing sequence such that

$$\lambda_m X_m = o(1) \text{ as } m \rightarrow \infty$$

$$\sum_{n=1}^m n X_n |\Delta^2 \lambda_n| = o(1)$$

and

$$\sum_{n=1}^m \frac{1}{P_n} |d_n|^k = o(X_m) \text{ as } m \rightarrow \infty,$$

where d_n is the n -th $(C, 1)$ transform of the sequence (na_n) then the series $\sum_{n=1}^{\infty} a_n \lambda_n$ is summable $|N, p_n|_k$, $k \geq 1$.

Putting $q_n = 1$ for all n , we get the corollary.

REFERENCES

- [1] Bor, H: On absolute summability factors, *Proceedings of the American Mathematical Society*, **118** (No.1) May 1993, pp. 71-75.
- [2] Borwein, D: On product of sequence, *Jour. Lond. Math. Soc.* **33** (1958) pp. 352-357.
- [3] Hardy, G. H : *Divergent Series*, Oxford Univ. Press, New York and London, 1949.
- [4] Mazhar, S.M: On $|C, 1|_k$ summability factors of infinite series, *Indian J. Math.* **14** (1972), pp. 44-45.
- [5] Mohanty, D: A study of some aspects of the theory of summability, *Ph.D. Thesis* (June, 1995), Berhampur University, Berhampur, Orissa, India, under the guidance of Prof. B.K. Tripathy.

Debadutta Mohanty

Department of Mathematics
Seemanta College, Jharpokharia
Mayurbhanja, Orissa, INDIA