

# Synthesis of some new Derivatives of N-[4-phenylthiazolyl]-3-Chloro- 4-[3'-iodo-4'-hydroxy-5'-methoxy]-2'-azetidinones with possible fungicidal activity

DR. JAGMOHAN SINGH NEGI

---

**ABSTRACT:** The present study was conducted to evaluate the fungicidal activity, many phenols and compounds with phenolic groups have antifungal potency. A large number of fungicides are formulated as wettable powders; this is the form most commonly used for spray mixes. They simply inhibit fungus growth temporarily. If the fungus is freed from such substance, it would revive. Such a chemical is called a "fungistat" and the phenomenon of temporarily inhibiting the growth is "fungistasis". Some other chemicals, like certain phenanthrene derivatives and Bordeaux mixture, may inhibit spore production without affecting the growth of vegetative fungistate hyphae. 2-amino 4-Phenyl thiazole is condensed with appropriate ethanol and piperidine aromatic was refluxed on water bath for 1 hr. Various obtaining gave benzal imine and azetidinones respectively and synthesized compounds showed moderate to good antifungal activity with respect to standard drugs.

**KEYWORDS:** 2-amino 4-Phenyl thiazole, EtOH, anhydrous zinc chloride, antifungal activity.

---

## 1. INTRODUCTION

The medicinal chemistry is the science which concerns essentially the understanding and explanation of mechanism of the action of drugs. For the chemist point of view it involves isolation, characterisation and synthesis of compounds, that can be used in medicines for the prevention, treatment and cure of diseases. Medicinal chemistry thus provides the chemical basis for the interdisciplinary field of therapeutics. The impact of the study of fundamental medicinal chemistry has given a new impetus to chemical, biological and engineering sciences. The approach to the study and design of the medicinal drug has centered primarily on the gross chemical structure of natural and synthetic compounds having established biological action. Modification of the basis structure are obtained by chemical synthesis and the effect of these changes on biological response are used to compile structure activity relationship. These relationship are intended to serve as a guide in the interpretation of the structural features essential for a given type of drug or new agents of similar biological activity. More commonly it is found that compounds are unrelated chemically may have the same action, and that compounds with the same functional group possess widely different biological activity. Such divergence from established structure activity relationship requires the synthesis of a large number of compounds before a useful agent can be found. A fungicide which kills on contact can check the growth of mycelium and limit the production of reproductive structures, such action delays or prevents the disease from spreading from infected plants to healthy ones near by, but the majority of successful non-systemic fungicides are most effective when applied prior to the arrival of infection. Non systemic fungicides must therefore normally possess some degree of persistence and majority of such fungicides are infected insoluble in water. In the present investigation synthesis of some new series Schiff bases derived from 2-Amino -4-


---

<sup>1</sup> Department of Chemistry, P.N.G. Govt P.G. College Ramnagar, Nainital, Uttarakhand (INDIA), E-mail address:- jagmohann011@gmail.com

phenyl thiazole with iodo vanillin is to be carried out which on cyclisation with corresponding thiazole derivative.

## MATERIAL AND METHODS

Thiazoles are important class of natural and synthetic compounds. Thiazole derivatives display a wide range of biological activities such as cardiotonic, fungicidal, sedative, anesthetic, bactericidal and anti-inflammatory. The synthesis of thiazole derivatives is important of their wide range of pharmaceutical and biological properties. A large number of fungicides are Fungicide which is effective only if applied prior to fungal infection is called a protectant; caused infection and thereby "curing" the plant, is called atherapeutant 8-quinolinol, antibiotics like Aureofungin, etc. Eradicants are those which remove pathogenic fungi from an infection court some chemicals do not kill fungi. The IR spectra were recorded on IR affinity-1, DRS-8000A, Shimadzu, Ptc. Ltd., Japan spectrophotometer. The <sup>1</sup>H-NMR was recorded in DMSO on Bruker Advance II 400 MHz spectrometer using TMS as an internal standard. Melting points were determined in open capillary tubes and are uncorrected. The purity of the compounds was checked by TLC-using Silica gel-G (Merck). Column chromatography was performed on silica gel. All the compounds were tested for their



2-Amino-4-(P-subst / Un-subst)-phenyl thiazole

## SYNTHESIS OF 2-AMINO -4- PHENYL THIAZOLE

A mixture of acetophenone (12.0 gm, 0.1 mol.) thiourea 15.2 gm.0.2 mol.) and iodine 25.4 gm.0.1 mol) was heated for 10 hrs on a steam bath. The crude reaction mixture are cooled and repeatedly extracted with ether to remove unreacted acetophenone and iodine. The residue was then dissolved in hot water & filtered to remove sulphur and other impurities. The solution was then moderately cooled and made alkaline with concentrated ammonia. 2-amino-4-phenyl thiazole, thus precipitated was collected and re-crystallised from dilute ethanol as long colourless, Yield : 65.5%, M.p: 149°C. % of N and S Calculated 16.27% and 37.20%, % of N and S found 16.20 and 36.20%. The compound has been characterized on the basis of I.R spectra and the purity of the compound was checked by T.L.C.

**[1] SYNTHESIS OF N-(4-PHENYL-2-THIAZOLYL)-2-IMINO-(3-IODO-4'-HYDROXY-5'-METHOXY BENZALIMINE) :**

A mixture of 2-Amino-4-phenyl thiazole (0.01 mol) and vanillin 0.01 moles in ethanol 30 ml and piperidine 3-4 drops was refluxed on water bath for 1 hours. the reaction mixture was cooled and the solid separated was filtered and recrystallised from ethanol. Yield: (92%), m.p. 160°C, IR(KBr) = 1210-1220 cm<sup>-1</sup> (due to C-O-c), 1665-1670 cm<sup>-1</sup> (C=N), 1590 - 1595 cm<sup>-1</sup> (C=C), 3000-3110 cm<sup>-1</sup> (due to -OH), 1640-1625 cm<sup>-1</sup> and 1250 cm<sup>-1</sup> (due to C=N and C-N), PMR = δ 4.0-4.2(3H, s, OCH<sub>3</sub>), δ 7.1-7.6(8H, m, ArH), δ 8.2-8.5(1H, s =CH), δ 9.5-.

**[2] SYNTHESIS OF N-[4-PHENYL-2-THIAZOLYL]-3-CHLORO-4-[3'-IODO-4'-HYDROXY-5-METHOXY)-2'-AZETIDINONES**

To a mixture of compound first (0.01 moles) and mercaptoacetic acid 0.001 mole dissolved in dioxane (20ml) a pinch of anhydrous zinc chloride was added and the mixture was refluxed for eight hours on cooling the separated solid was washed with dilute sodium bicarbonate was crystallized from ethanol. Yield 47, M.P 148°C IR(KBr) = 3100 cm<sup>-1</sup> (due to OH), 1640-1625 cm<sup>-1</sup> AND 1250 (C=N), 1590 - 1595 cm<sup>-1</sup> (C=C), 3000-3110 cm<sup>-1</sup> (due to -OH), 1640-1625 cm<sup>-1</sup> and 1250 cm<sup>-1</sup> (due to C=N and C-N), 1210-1220 CM<sup>-1</sup> (due to C-O-C), 1660-1670 cm<sup>-1</sup> (due to C-S-C) 1685 cm<sup>-1</sup> (due to cyclic > c=O), PMR = δ 3.82-3.86 (3H, s, OCH<sub>3</sub>), δ 9.85(1H, s, OHm), δ 4.15-(2H, s, CH<sub>2</sub>S), δ 6.5-6.8(1H, s, -CH), δ 6.5-7.4(8H, M, -Ar-H), Similarly, various N-[4-(P-subst/un-subst)-Phenyl-2-thiazolyl]-2-imino-(3'-iodo-4'-Hydroxy-3'-methoxyphenyl)-4'-thiazlidone were prepared by using similar reaction procedure and their analytical data are incorporated in the table (II) respectively antibacterial activity against *S. pyogenus*, while rest of all derivatives possessed good activity against *S. pyogenus* in the range of 125-250 µg/ml. While rest of all derivatives are poor against *A. Niger*.

**TABLE I : Analytical data of subst/un-subst)-Phenyl-2-thiazolyl]-2-imino-(3-iodo-4'-Hydroxy-3'-methoxy benzyldine.**

S.N.	Nature of Ar	Molecular Formula	Yield %	M.P. °C	ELEMENTAL ANALYSIS			
					% of N		% of S	
					Cald	Fond	Cald	Found
la	2-Amino-4-phenyl thiazole	C <sub>17</sub> H <sub>13</sub> N <sub>2</sub> O <sub>2</sub> Sl	42	138	9.03	09.00	10.32	10.25
lb	2-Amino-4(p-chloro)-phenyl thiazole	C <sub>17</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub> ClSl	50	140	19.92	19.86	22.77	22.69
lc	2-Amino-4(p-fluoro)-phenyl thiazole	C <sub>17</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub> SFl	52	145	08.53	08.50	09.75	09.70
ld	2-Amino-4(p-bromo)-phenyl thiazole	C <sub>17</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub> SBrI	48	106	07.21	07.11	08.24	08.20
le	2-Amino-4(p-nitro)-phenyl thiazole	C <sub>17</sub> H <sub>12</sub> N <sub>3</sub> O <sub>4</sub> Sl	47	148	11.83	11.76	09.01	08.93
lf	2-Amino-4(p-hydroxy)-phenyl thiazole	C <sub>17</sub> H <sub>15</sub> N <sub>2</sub> O <sub>3</sub> Sl	48	165	08.53	08.49	09.75	09.73
lg	2-Amino-4(p-methyl)-phenyl thiazole	C <sub>20</sub> H <sub>16</sub> N <sub>2</sub> O <sub>3</sub> Sl	52	226	08.64	08.60	09.87	09.80
lh	2-Amino-4(p-methoxy)-phenyl thiazole	C <sub>17</sub> H <sub>15</sub> N <sub>2</sub> O <sub>4</sub> Sl	53	246	08.23	08.20	09.41	09.30
li	2-Amino-4(p-ethoxy)-phenyl thiazole	C <sub>20</sub> H <sub>16</sub> N <sub>2</sub> O <sub>3</sub> Sl	50	250	07.90	07.80	09.03	08.90

**TABLE-2 : Analytical data of N-[4-(p-subst/un-subst)Phenyl-2-thiazolyl]-3-chloro-4-[3'-iodo-4'-hydroxy-5-methoxy)-2'-azetidinone**

S.N.	Nature of Ar	Molecular Formula	Yield %	M.P. °C	ELEMENTAL ANALYSIS			
					% of N		% of S	
					Cald	Fond	Cald	Found
lla	2-Amino-4-phenyl thiazole	C <sub>19</sub> H <sub>13</sub> N <sub>2</sub> O <sub>3</sub> Cl I	52	179	07.09	07.05	16.24	16.22
llb	2-Amino-4(p-chloro)-phenyl thiazole	C <sub>19</sub> H <sub>12</sub> N <sub>2</sub> O <sub>3</sub> SCl <sub>2</sub> I	53	190	06.50	06.45	14.93	14.90
llc	2-Amino-4(p-fluoro)-phenyl thiazole	C <sub>19</sub> H <sub>12</sub> N <sub>2</sub> O <sub>3</sub> SFCI I	50	185	06.54	06.50	15.05	15.00
lld	2-Amino-4(p-bromo)-phenyl thiazole	C <sub>19</sub> H <sub>12</sub> N <sub>2</sub> O <sub>3</sub> SBrCl I	42	186	05.93	05.88	13.55	13.52
lle	2-Amino-4(p-nitro)-phenyl thiazole	C <sub>19</sub> H <sub>12</sub> N <sub>3</sub> O <sub>5</sub> SCl I	50	225	09.56	09.49	14.97	14.55
llf	2-Amino-4(p-hydroxy)-phenyl thiazole	C <sub>20</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub> S <sub>2</sub> Cl I	42	148	06.82	06.78	15.60	15.55
llg	2-Amino-4(p-methyl)-phenyl thiazole	C <sub>20</sub> H <sub>15</sub> N <sub>2</sub> O <sub>4</sub> Cl I	51	144	06.86	06.81	15.68	15.60
llh	2-Amino-4(p-methoxy)-phenyl thiazole	C <sub>20</sub> H <sub>15</sub> N <sub>2</sub> O <sub>5</sub> Cl I	50	141	06.60	06.55	15.09	14.55
lli	2-Amino-4(p-ethoxy)-phenyl thiazole	C <sub>21</sub> H <sub>17</sub> N <sub>2</sub> O <sub>4</sub> SCl I	54	145	06.39	06.81	14.61	14.02

## ANTIFUNGAL SCREENING

The newly synthesized compounds were evaluated against *Alternaria alternate* fungus at optimum temperature of  $28 \pm 1^\circ\text{C}$  (after 7 days incubation) was observed. After inoculation, All the petridishes were incubated at  $(25 \pm 20^\circ\text{C})$  for 7 days, the efficiency of various ant-fungal was recorded by measuring the radial growth of the fungal colony (in mm). The percentage inhibition of fungus mycelia growth was calculated by the equation.


$$\% \text{ of Inhibition} = \frac{|(C - T) \times 100|}{C}$$

Where C and T are average colony diameters (in mm) of the fungal colony in control © and treated (T) plates respectively.

**Table-3-Effect of Some Newly Synthesised Antifungal Compounds against *Fusarium solani* at optimum temperature (After 7 days incubation)**

Compound	Dose	Average colony diameter (in mm) in PDA medium	% Inhibition
Control		60.88	
la	0.20	2.7	95.38
lb	0.20	3.2	94.73
lc	0.20	4.0	93.42
ld	0.20	1.9	96.87
le	0.20	2.7	95.55
lf	0.20	2.7	95.38
lg	0.20	9.9	83.71
lh	0.20	3.0	95.06
li	0.20	3.2	94.73
lla	0.20	3.1	94.90
llb	0.20	2.7	95.55
llc	0.20	4.1	93.25
lld	0.20	3.5	94.24
lle	0.20	3.2	94.73
llf	0.20	2.8	95.39
llg	0.20	2.4	96.05
llh	0.20	1.7	97.20
lli	0.20	2.8	95.39
BAVISTIN(Std drug)	0.20	0.22	99.65





## CONCLUSION

It is evident from fungal screening data that all the newly synthesized compound tested were found satisfactorially superior over control but inferior to that of standard antifungal (Bavistin) compound. Mostly synthesized compound showed marked of the fungal growth in vitro test. It can also be concluded from the result that mostly synthesized compound are good antifungal and showed significant level of antifungal activity and

## RESULTS AND DISCUSSIONS

It is evident from fungal screening data that all the newly synthesized compound tested were found satisfactorially superior over control but inferior to that of standard antifungal (Bavistin) compound. Mostly synthesized compound showed marked of the fungal growth in vitro test. It can also be concluded from the result that mostly synthesized compound are good antifungal and showed significant level of antifungal activity and compound No (lg) showed moderate activity.

## ACKNOWLEDGEMENTS

The authors are thankful to the Dr. Adarsh Pandey, Head Department of Botany, G.F.P.G. College Shahjahanpur for providing necessary facilities the antimicrobial activity and also thankful to CDRI Lucknow for providing IR spectra and micro-care laboratory

## REFERENCES

1. Amr, Ael-G, Abdel-Lalif, N. A., Abdalla, M. M., Bioorg Med. Chem., 2006, 14(2), 373
2. Anandarajagopal, K., Sunilson A. J., Illavarasu, A. Thangavelpandian, N., R., Int. J. ChemTech. Res., 2010, 2(1), 45.
3. Fu, L., Feng, X., Wang, J.J., Xun, Z., Hu, J.D., Zhang, J.J., Zhao, Y.W., Huang, Z.B., Shi, D.Q. Efficient synthesis and evaluation of antitumor activities of novel functionalized 1,8-naphthyridine derivatives, *Comb. Sci.*, **2015**, 17, 24–31.
4. Ashok, K., Rathod, IDiseant. J. PharmTech. Res. 2011, 3(1), 435.
5. A.L. Barry, "The antimicrobial. suceptibility Test Principal and Practices, . Edicted by Illus lea and Fabiger 180, Bio. . Abstr. 64 (1976). 25183
6. Abid, M., Azam, A., Bioorg. Med. Chem. Lett., . 2006, 16(10), 2812
7. Al-Saddi. M. S; Faidallah. H. M; Rostom.S.A.F. Arch. Pharm. Che. Life Sci., 2008, 341, 9.42
8. B.Das and M.k.Rout. J Indian. Chem.Soc. 1955. 32. 668.
9. Tripathi. K. D. Essential of medical pharmacology., 2003, 5th edition, 627- 686
10. Karpov K. A; Nazarenko A. V; Pekarevskii B. V; potekhin. V. M. Russian journal of applied chemist 2001, 74, 998-1001
11. De los Rios, C., Egea, J., Marco-Contelles, J., Leon, R., Samadi, A., Iriepa, I., Moraleda, I., Galvez, E., García, A.G., Lopez, M.G., Villarraya, M., Romero, A. Synthesis, inhibitory activity of cholinesterases, and neuroprotective profile of novel 1,8-naphthyridine derivatives, *J. Med. Chem.*, **2010**, 53, 5129–5143.
12. Baselt. T; Rehse. K. Archived der pharmazie., 2008, 24, 645- 654
13. Ashok, K., Rathod, IDiseant. J PharmTech. Res. **2011**, 3(1), 435.

14. Abid, M., Azam, A., Bioorg. Med. Chem. Lett., **2006**, 16(10), 2812
15. Amr, Ael-G, Abdel-Lalif, N. A., Abdalla, M. M., Bioorg. Med. Chem., 2006, 14(2), 373
16. Anandarajagopal, K., Sunilson, A. J., . ss Illavarasu, A., Thangavelpandian, N., R., Int. J. ChemTech. Res., **2010**, 2(1), 45
17. Hansford, K. A., Zanzarova, V.Dorr, . A., Lubell, W. D., JComb. Chem. **2004**, 6, 893.
18. Heckmann, G, Bach, T., Angew. Chem. Int. Ed. **2005**, 44, 1199. Millan, S. E Nicholas, D. P., . Bioorg. Med. Chem. Lett., **2009**, 19, 222.
19. Krishna, R., Pande, B. R., Bharthwal, S. P., Parmar, S. S., Eur. J. Med. Chem., **1980**, 15, 567 .
20. Lester A., Itschep M. and Foye, Principle of Medicinal Chemistry, **5**, 833 (**2002**).
21. Regaila, H. A., El-Bayonk, A. K., Hammad, M., Egypt J. . Chem., **1979**, 20, 19
22. Sharma H.L., Principle of Pharmacology, **2**, 748-752 (**2008**)
23. Schimdt, P., Eichenberger, K., Schweizer, E., Germa Offen.196908479, Chem , Abstr. **1970**, 72, 31837u .
24. Satoskar R.S. and Bhandarkar S.D., Pharmacology and Pharmacotherapeutics, **2**, (**2007**)
25. Fu, L., Lin, W., Hu, M.H., Liu, X.C., Huang, Z.B., Shi, D.Q. Efficient synthesis of functionalized benzo[*b*] [1,8]naphthyridine derivatives via three-component reaction catalyzed by l-proline, *ACS Comb. Sci.*, **2014**, 16, 238–243.
26. Sampathkumar, N., Muruges, A., Rajendran, S.P. Afacile synthesis of 5-phenyl-dibenzo[*b*, *g*][1,8]naphthyridines, *J. Heterocycl. Chem.*, **2015**, 53, 924–928.
27. Zhang, H.M., Dai, H., Hanson, P.J., Li, H., Guo, H., Ye, X., Hemida, M.G, Wang, L., Tong, Y., Qiu, Y., Liu, S., Wang, F., Song, F., Zhang, B., Wang, J.G, Zhang, L.X., Yank, D. Antiviral activity of an isatin derivative via induction of PERK-Nrf2-mediated suppression of cap-independent translation, *ACS Chem. Biol.*, **2014**, 9, 1015–1024.





This document was created with the Win2PDF “print to PDF” printer available at <http://www.win2pdf.com>

This version of Win2PDF 10 is for evaluation and non-commercial use only.

This page will not be added after purchasing Win2PDF.

<http://www.win2pdf.com/purchase/>