

## Evaluation of Pre and Post Emergence Herbicides In Chickpea (*Cicer arietinum*L.)

A. K. Gore\*, S. M. Gobade\* and P.V. Patil

**ABSTRACT:** An agronomic investigation on "Evaluation of pre and post emergence herbicides in irrigated chickpea (*Cicer arietinum* L)" was carried out on research farm of Agronomy Department, Vasantrya Naik Marathwada Krishi Vidyapeeth, Parbhani during Rabi 2011-12. Results revealed that Weed free (weeding at 20 days interval up to 80-90DAS) recorded highest weed control efficiency, grain, straw and biological yield. It was found at par with of Pendimethalin (PE) @ 0.75 a.i. kg ha<sup>-1</sup>+ 1HW, Pendimethalin (PE) @ 1.00 a.i. kg ha<sup>-1</sup>, 1 Hoeing (30 DAS) + (2 hand weedings) and Mechanical weedings (2 hand weedings) and significantly superior over rest of the treatments. Among the chemical weed control treatment application of Pendimethalin (PE) @ 0.75 a.i. kg ha<sup>-1</sup>+ 1 HW was found beneficial higher grain, straw yield, weed control efficiency and effective in controlling weeds and increasing the yield of chickpea.

**Keywords:** Chickpea, Herbicide, Pre-emergence, Post-emergence.

### INTRODUCTION

Chickpea is an important pulse crop of the semi-arid tropics, particularly in the rainfed ecology of the Indian subcontinent. In the last decade, this crop has experienced an export-driven expansion in new riches such as Australia and Canada. Globally, chickpea is cultivated on about 10.4 million ha area adding 8.57 million tonnes of seeds to the global food market, with an average productivity of 826 kg ha<sup>-1</sup>. As many as 45 countries grow chickpea. India grows chickpea on about 8.56 million ha producing 7.35 million tonnes seeds with productivity 858 kg ha<sup>-1</sup> which represents 37 per cent and 50 per cent of the national pulse acreage and production respectively. The important chickpea growing states are Madhya Pradesh, Rajasthan, Uttar Pradesh, Maharashtra, Karnataka and Haryana. In Maharashtra state the area was 13.95 lack ha and production was 13.01 lack tonnes with productivity of 933 kg ha<sup>-1</sup> during the year 2011-12. (Anonymous., 2011).

The average yield of this crop is very much which may be due to many factors but among these infestation of weeds is very important. (Surjit Singh *et al*, 2008). Chickpea, being slow in its early growth and short stature plant, is highly susceptible to weed competition and often considerable losses may occur if weeds are not controlled at proper time. Weeds are

serious constraint to increase production and easy harvesting in chickpea. Yield losses as due to weeds were observed to vary between 40-90 per cent. *Chenopodium album* also one of the dominant weed in chickpea and causes maximum reduction in grain yield. (Mailk and Balyan., 1988). Moreover potential yield losses in chickpea due to weeds range between 22 to 100 per cent. Weeds reduce grain yield of chickpea up to an extent of 60 per cent (IIPR., 2004).

Hand weeding is practiced in traditional production areas, but is impractical in the impractical in the extensive production areas. Hand weeding is labour-intensive and therefore an expensive operation when done by hired labour and, if delayed, the operation does not prevent adverse effect of the weeds on crop yield. The use of appropriate herbicides can eliminate this early weed competition and prevent yield losses Herbicides are selective, cost effective, easy to apply, and offer flexibility in application time. (Peterson *et al.*, 2001 or Hoseinyet *al.*, 2011). When properly used, pre-emergence herbicides accomplish effective and economic weed control, and consequently chickpea seed yields as similar to or only slightly smaller than those of weed free treatments are resulted (Gul Hassan *et al.*, 2007). New pre and post emergence weedicides are available in market for effective control of weeds. It is therefore felt

\* Vasantrya Naik Marathwada Agriculture University, Parbhani 431 402, E-mail : akgoremko@gmail.com., gobadeagricos@gmail.com

necessary to study the efficacy of these new weedicides for control of weeds in gram and also to compare these herbicides with the exiting cultural weed control methods.

## MATERIAL AND METHODS

An agronomic investigation on "Evaluation of pre and post emergence herbicides in irrigated chickpea (*Cicerarietinum* L.) on black cotton soils of marathwada region." was carried out on research farm of Agronomy Department, Vasanttrao Naik Marathwada Krishi Vidyapeeth, Parbhani during *Rabi* 2011-12 to evaluate pre and post emergence herbicides in irrigated chickpea for effective and economical weed control in chickpea. The experiment was conducted on vertisols, in a randomized block design with twelve treatments four pre emergence herbicides viz, Pendimethalin (PE) @ 0.75 a.i kg ha<sup>-1</sup>+ 1 HW, Pendimethalin (PE) @ 1.00 a.i kg ha<sup>-1</sup> Trifluralin (PE) @ 1.0 a.i. kg ha<sup>-1</sup>, Oxyfluorfen @ 0.125 a.i. kg ha<sup>-1</sup>, and Metribuzin @ 0.75 a.i. kg ha<sup>-1</sup> and three post-emergence herbicides viz, Imazethapyr (POE) @ 0.75 a.i. kg ha<sup>-1</sup>, Quizalofop-p-ethyl (POE) @ 40 a.i. g ha<sup>-1</sup>, Propaquizafop (POE) @ 0.75 a.i. kg ha<sup>-1</sup> and three cultural treatments viz, 1 Hoeing (30 DAS) + (2 hand weedings), Mechanical weedings (2 hand weedings), Weed free (weeding at 20 days for up to 80-90 DAS), and Weedy check. They were replicated three times. The gross and net plot sizes were 5.4 m x 4.5 m and 4.5 m x 3.6 m, respectively. The seeds of variety Vijay were sown by dibbling method at spacing 45 cm x 10 cm. Observations on growth, weed parameter were

recorded periodically at an interval of 30 days. Whereas the observation on yield attributing characters and yield were recorded at harvest.

## RESULT AND DISCUSSION

Data in respect of grain yield (kg ha<sup>-1</sup>), straw yield (kg ha<sup>-1</sup>), biological yield (kg ha<sup>-1</sup>), and weed control efficiency (%) of chickpea as influenced by various treatments are presented in Table 1.

Data furnished in Table 1 indicate that seed yield, straw yield and biological yield of chickpea was significantly influenced by various weed control treatments.

Treatment weed free (weeding at 20 days interval up to 80-90 DAS) recorded significantly higher seed yield (2976 kg ha<sup>-1</sup>), straw yield (3247 kg ha<sup>-1</sup>) and biological yield (6223 kg ha<sup>-1</sup>) of chickpea and it was found at par with 1 Hoeing (30 DAS) + (2 HWs), Mechanical weedings (2 hand weedings), and Pendimethalin (PE) @ 0.75 kg ha<sup>-1</sup>. Whereas it was found significantly superior over rest of the weed control treatments and weedy check. Thus the effective weed control achieved in the earlier mentioned treatments resulted in enhancing various growth and yield attributing characters of chickpea and finally gave significantly higher grain and straw yield over Weedy check. Low yield in Weedy check may be due to poor root growth and higher weed population could have competed with chickpea crop for space, water and nutrients, there by adversely affecting grain and straw yield, Similar trend was observed Pooniya *et al.* (2009) and Ratnum *et al.* (2011).

**Table 1**  
Grain yield, straw yield, biological yield kg ha<sup>-1</sup> and weed control efficiency of chickpea as influenced by various treatments

Treatments	Grain Yield (Kg ha <sup>-1</sup> )	Straw Yield (Kg ha <sup>-1</sup> )	Biological Yield (Kg ha <sup>-1</sup> )	Weed control efficiency (%)	
				Monocot (At harvest)	Dicot weed (At harvest)
T <sub>1</sub> : Pendimethalin (PE) @ 0.75 a.i. kg ha <sup>-1</sup>	2796	3048	5844	69.76	76.14
T <sub>2</sub> : Trifluralin (PE) @ 1.0 a.i. kg ha <sup>-1</sup>	2450	2726	5176	63.67	75.73
T <sub>3</sub> : Metribuzin (PE) @ 0.75 a.i. kg ha <sup>-1</sup>	2010	2413	4423	73.24	75.44
T <sub>4</sub> : Oxyfluorfen (PE) @ 0.125 a.i. kg ha <sup>-1</sup>	2223	2516	4739	68.02	74.81
T <sub>5</sub> : Mechanical Weedings (2 hand weedings)	2813	3083	5896	74.71	76.89
T <sub>6</sub> : Imazethapyr (POE) @ 0.75 a.i. kg ha <sup>-1</sup>	2153	2490	4643	74.44	74.4
T <sub>7</sub> : Quizalofop-p-ethyl (POE) @ 40 a.i. g ha <sup>-1</sup>	2435	2719	5154	76.98	73.19
T <sub>8</sub> : Propaquizafop (POE) @ 0.75 a.i. kg ha <sup>-1</sup>	2394	2683	5077	76.92	50.89
T <sub>9</sub> : 1 Hoeing (30DAS) + (2 hand weedings)	2950	3233	6183	78.72	82.09
T <sub>10</sub> : Weed free (weeding at 20 days interval up to 80-90 DAS)	2976	3247	6223	85.35	85.09
T <sub>11</sub> : Weedy check	1540	1870	3410	0	0
S.E. ±	137	175	336	—	—
C.D. at 5 %	412	525	1008	—	—
GM	2440	2739	5160	74.14	74.46




Figure 1: Grain yield kg ha<sup>-1</sup>, Bhoosa yield kg ha<sup>-1</sup>, and Biological yield kg ha<sup>-1</sup> as influenced by various treatments




Figure 2: Monocot weed control efficiency (%) as influenced periodically by various treatments




Figure 3: Dicot weed control efficiency (%) as influenced periodically by various treatments

As we see the effect of chemical weed control methods, the application of Pendimethalin (PE) @ 0.75 kg ha<sup>-1</sup> recorded significantly higher grain yield (2796 Kg ha<sup>-1</sup>), straw yield (3048 Kg ha<sup>-1</sup>) and biological yield (5844 Kg ha<sup>-1</sup>) which was found at par with Trifluralin (PE) @ 40 g ha<sup>-1</sup> and Propaquizafop (PE) @ 0.75 kg ha<sup>-1</sup> and it was found significantly superior over rest of chemical weed control treatment and weedy check. Similar trend was observed Pooniya *et al.* (2009) and Ratnum *et al.* (2011).

In case of weed control efficiency of chickpea generally among all the weed control treatments weed free recorded highest weed control efficiency at all the growth stages for dicot as well as monocot weeds. Among herbicidal treatments it was highest in Pendimethalin (PE) @ 0.75 kg ha<sup>-1</sup> for dicot weeds and in Quizalofop-p-ethyl (POE) @ 40 g ha<sup>-1</sup> for monocot weeds at all the stages of observations. Similar finding observed by Sharma *et al.* (2006).

## REFERENCES

- Anonymous (2011), Chemical weed control in chickpea (*Cicerarietinum*L.) submitted thesis dissertation at MKV, Parbhani.
- Gul Hasan and Imtiaz Khan (2007), Post emergence herbicidal control of *Asphodelustenuifolius* in desi chickpea (*Cicerarietinum* L.). *Pakistan J. weed sci. Res* **13** (1-2): 33-38.
- Hoseiny-Rad., Manijeh, Jagannath and Shobha (2011), Effect of herbicide Imazethapyr (pursuit) on chickpea seed germination. *Archives of Phytopathology and Plant Protection*, **44**(3): 224-230(7).
- IIPR, (2004), *Annual Report*. Indian Institute of Pulses Research, Kanpur. pp. 18-19.
- Malik, R. K. and Balyan, R. S. (1988), Effect of herbicides and planting dates in chickpea. *Indian Journal of Weed Sci.* **20** (2): 75-81.
- Pooniya Vijay, B. Rai and R.K. Jat (2009), Yield and yield attributes of chickpea (*Cicerarietinum* L.) as influenced by various row spacing and weed control. *Indian. J. weed sci.* **41** (3 & 4): 222-223.
- Sharma, R. K. and V. K. Goswami (2006), Integrated weed management in chickpea (*Cicerarietinum* L.). *Indian J. weed Sci.* **20**(2), 102-103.
- Surjit Singh., U.S. Walia and Buta Singh (2008), Effective control of weeds in chickpea. *Indian J. Weed. Sci.* **40** (1 & 2): 31-55.