



INTERNATIONAL JOURNAL OF TROPICAL AGRICULTURE

ISSN : 0254-8755

available at <http://www.serialsjournal.com>

© Serials Publications Pvt. Ltd.

Volume 35 • Number 3 • 2017

Effect of Zinc Solubilizing Microorganisms on Yield and Economics of Groundnut Grown on Vertisol

Prasad Sable^{1*}, Syed Ismail² and Sonwane N.S.³

¹ M.Sc. Scholar, Department of Soil Science and Agricultural Chemistry, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani (MS)-431402

² Associate Professor, Department of Soil Science and Agricultural Chemistry, Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani (MS)-431402

³ Environmental Scientist, SPMESM, Aurangabad (MS)-431006

* E-mail: sable.prasad7@gmail.com

Abstract: The experiment was conducted on zinc deficient soil at farmer's field in Parbhani, Maharashtra during summer season of 2014-15 to assess the ability of zinc solubilizing microorganisms on yield and economics of Groundnut. The treatments comprised of seven strains of zinc solubilizers viz. *Burkholderia cepacia*, *Burkholderia cenocepacia*, *Pseudomonas fluorescens*, *Pseudomonas striata*, *Trichoderma viride*, *Trichoderma harzianum* and *Bacillus megaterium* along with recommended dose of fertilizers which were replicated three times in randomized block design. The result revealed that kernel and haulm yield of groundnut and net returns was increased with inoculation of zinc solubilizing microorganisms along with recommended dose of fertilizers over uninoculated control. Among all the zinc solubilizers, RDF + *Rhizobium* + *Pseudomonas striata* recorded significantly highest yield and net returns.

Key words: Groundnut, GMR, NMR, zinc solubilizing microorganism.

INTRODUCTION

The Oil seed crops have been the backbone of agricultural economy of India and the world. Groundnut (*Arachis hypogaea* L.) is an important oil seed crop and food grain legume. The groundnut

has indeterminate growth habit; hence growth and development of reproductive and vegetative organs overlap. This causes low fruiting efficiency due to inter organ competition for photo-assimilates and other metabolites. Consequently there is improper

partitioning of assimilates to the developing pods and seeds. Most prominent constraint in the low yield is extended duration of flowering and variable pod sizes. Parmar and Schmidt (1989) have demonstrated the translocation of photosynthates within groundnut plant is not random but has a definite pattern, and this pattern is changed during different phases of plant growth. In view of the vagaries of rainfall and soil fertility problems, groundnut crop needs efficient nutrient management practices with an integrated approach of exploring all possible nutrient resources, not only to augment the nutrient supply but also to improve the physical, chemical and biological properties of soil, so as to overcome the limitations of groundnut cultivation.

At several places normal yield of crops could not be achieved despite judicious use of NPK fertilizers due to deficiency of micronutrients in soil in general that of Zn in particular. Numerous microorganisms especially those associated with roots have the ability to increase plant growth and productivity. In soil, both macro and micronutrients undergo a complex dynamic equilibrium of solubilization and insolubilization that is greatly influenced by the soil pH and micro flora and that ultimately affects their accessibility to plant roots for absorption (Altomare *et al.*, 1999). Keeping these points in consideration, field trial was conducted to enhance the yield and economics of groundnut using Zn solubilizing microbial cultures on Vertisol.

MATERIALS AND METHOD

The experiment was conducted on farmer's field in Kehal village, Tq. Jintoor, Dist. Parbhani on Vertisol during summer season of 2014-15 on Groundnut. The soil was clayey in texture, moderately alkaline in reaction, medium in available nitrogen, phosphorus and sufficient in available potassium, iron, sulphur and low in zinc. As among the micronutrient status Cu, Fe and Mn were above the critical limits but Zn content in experimental soil was found to be

deficient. The treatments comprising inoculation with liquid inoculants of *Bradyrhizobium* and *Bacillus megaterium* (PSB) for Groundnut seed in alone and in combinations. Total eight treatments of bioinoculants were replicated three times in RBD. The experiment consists of 8 treatments of laboratory tested P and Zn solubilizers RDF+ *Rhizobium*; RDF + *Rhizobium* + *Bacillus megaterium*; RDF + *Rhizobium* + *Burkholderia cepacia*; RDF + *Rhizobium* + *Burkholderia cenocepacia*; RDF + *Rhizobium* + *Pseudomonas fluorescens*; RDF + *Rhizobium* + *Pseudomonas striata*; RDF + *Rhizobium* + *Trichoderma viride*; RDF + *Rhizobium* + *Trichoderma harzianum*. Seed treatment was done before sowing with liquid bioinoculants each @ 50 ml 10 kg⁻¹ seed. The crop was raised following recommended agronomic practices. The recommended dose of chemical fertilizers was applied @ 30:60:30 NPK kg ha⁻¹ at the time of sowing. Intercultural operations like thinning, weeding, spraying of insecticides, fertilizer application and schedule of irrigation for groundnut crop was carefully followed. The crop variety used was TAG 37A. The data obtained was statistically analyzed and appropriately interpreted as per the methods described in "Statistical Methods for Agricultural Workers" by Panse and Sukhatme, (1985).

RESULTS AND DISCUSSION

Yield of Groundnut

The results narrated in Table 1 indicate significant impact of Zn solubilizing microorganisms on dry pod and dry haulm yield of groundnut. Significantly highest dry pod yield and dry haulm yield of groundnut was noted in RDF + *Rhizobium* + *Pseudomonas striata* but treatment RDF + *Rhizobium* + *Pseudomonas fluorescens*, RDF + *Rhizobium* + *Trichoderma viride*, RDF + *Rhizobium* + *Trichoderma harzianum* and RDF + *Rhizobium* + *Burkholderia cenocepacia* were found statistically at par. Lowest biological yields were obtained in treatment T₁. These results are also corroborate with reports of Datta

Table 1
Effect of zinc solubilizing micro-organism on dry pod and dry haulm yield of Groundnut

Sr. No.	Treatment	Dry Pod yield (q ha ⁻¹)	Dry Haulm yield (q ha ⁻¹)
T ₁	RDF + <i>Rhizobium</i>	23.50	30.73
T ₂	T1 + <i>Bacillus megaterium</i>	24.96	32.19
T ₃	T1 + <i>Burkholderia cepacia</i>	25.63	32.86
T ₄	T1 + <i>Burkholderia cenocepacia</i>	26.67	33.90
T ₅	T1 + <i>Pseudomonas fluorescens</i>	28.43	35.66
T ₆	T1 + <i>Pseudomonas striata</i>	29.10	36.33
T ₇	T1 + <i>Trichoderma viride</i>	28.60	35.83
T ₈	T1 + <i>Trichoderma harzianum</i>	27.83	35.06
	S.E. ±	0.62	0.62
	C.D. at 5 %	1.88	2.61
	C.V. %	4.01	3.15

et al. (2014) as they revealed that pod and haulm yield of groundnut was significantly influenced by different nutrient management, *Rhizobium* inoculation of groundnut seed along with soil application of 50 kg P₂O₅ ha⁻¹ and 50 kg K₂O ha⁻¹. This may be owing

to the fact that seed inoculation with biofertilizer, soil phosphorus and potassium significantly increased the yield component and yield of groundnut. Thereafter, Gupta (2006) noted that the seed inoculation with *Rhizobium* + PSB + *Azotobacter* increased grain yield in chickpea. However, Sable *et al.* (1998) also reported significant increase in grain yield with inoculation with *Rhizobium* + RDF.

Economics of Groundnut

The data presented in the Table 2 revealed that gross monetary returns and net monetary return of groundnut was obtained significantly highest in treatment RDF + *Rhizobium* + *Pseudomonas striata* followed by treatments RDF + *Rhizobium* + *Pseudomonas fluorescens* and RDF + *Rhizobium* + *Trichoderma harzianum*. The treatment RDF + *Rhizobium* + *Pseudomonas fluorescens*, RDF + *Rhizobium* + *Trichoderma viride* and RDF + *Rhizobium* + *Trichoderma harzianum* were found to be at par with each other. Regarding, benefit cost ratio of groundnut, the highest B:C ratio was computed in treatment RDF + *Rhizobium* + *Pseudomonas striata* and T₅, T₇ and T₈ these treatments were found to be at par with each other. These results are concurred with Subrahmaniyan

Table 2
Effect of zinc solubilizing microorganisms on economics of groundnut

Sr. No.	Treatment	Cost of cultivation (Rs ha ⁻¹)	Gross monetary return (Rs ha ⁻¹)	Net monetary return (Rs ha ⁻¹)	B:C
T ₁	RDF + <i>Rhizobium</i>	30159	84600.00	54441.00	2.81
T ₂	T1 + <i>Bacillus megaterium</i>	30359	89856.00	59497.00	2.96
T ₃	T1 + <i>Burkholderia cepacia</i>	30359	92268.00	61909.00	3.04
T ₄	T1 + <i>Burkholderia cenocepacia</i>	30359	96012.00	65653.00	3.16
T ₅	T1 + <i>Pseudomonas fluorescens</i>	30359	102360.00	72001.00	3.37
T ₆	T1 + <i>Pseudomonas striata</i>	30359	104760.00	74401.00	3.45
T ₇	T1 + <i>Trichoderma viride</i>	30359	102960.00	72601.00	3.39
T ₈	T1 + <i>Trichoderma harzianum</i>	30359	100188.00	69829.00	3.30
	S.E. ±		2236.2	2236.2	0.07
	C.D. at 5 %	–	6783.5	6783.5	0.22
	C.V. %	–	4.01	5.84	4.01

et al., (1998) reported that highest benefit: cost ratio (B:C) was recorded in plot inoculated with *Rhizobium* culture and *Rhizobium* culture + 50 kg P₂O₅+ 50 kg K₂O ha⁻¹ compared to all other treatment. These results are also in close conformity with reports of Subrahmaniyan *et al.*, (2000) and Mohapatra *et al.*, (2010). Further, Naik *et al.*, (2012) reported that organic manure to safflower had significant influence on the economics of safflower production and also application of 150% NPK recorded significantly higher gross returns, net returns, cost of cultivation and B:C ratio over 100% NPK, but was on par with 125% NPK in black clay loam soil.

CONCLUSION

The inoculation of zinc solubilizing microorganisms along with NPK increased the yield of groundnut. Results revealed that significantly highest kernel and haulm yield of groundnut was noted in inoculated plots with liquid form of *Rhizobium* and *Pseudomonas striata* over other treatments. The significant improvement in economics (GMR, NMR and B:C) of groundnut crop was also noted with inoculation of *Rhizobium* and *Pseudomonas striata* along with recommended dose of fertilizers.

REFERENCES

- Altomare, C., Norvell, W. A., Bjorkman, T. and Harman, G. E. (1999), Solubilization of Phosphates and Micronutrients by the Plant-Growth-Promoting and Biocontrol Fungus *Trichoderma harzianum* Rifai 1295-22. *Applied And Environmental Microbiology* **65**(7): 2926–2933.
- Datta, M., Yadav, G.S, and Chakraborty, S. (2014), Integrated nutrient management in groundnut in tropical humid climate of north east India. *Indian Journal of Agronomy* **59**(2): 322-326.
- Gupta S.C. (2006), Effect of combined inoculation on nodulation, nutrient uptake and yield of Gupta S.C. (2006). Effect of combined inoculation on nodulation, nutrient uptake and yield of chickpea in Vertisols. *Journal of Indian Society of Soil Science*, **54**(2): 251-254.
- Mohapatra, A.K.B., and Dixit, L., (2010), Integrated nutrient management in rainy season groundnut. *Indian Journal of Agronomy* **55**(2): 123-127.
- Naik, R., Halepeyati, A. S., Desai, B. K. and Pujari, B. T. (2012), Influence of organic manure and nutrient levels on growth, yield components and yield of safflower (*Carthamus tinctorius* L.). *Journal of Oilseed Research*, (29, Spl. Issue): 257-258.
- Panase, U.G. and Sukhatme, P.V. (1985), Statistical Methods for Agricultural Workers. I.C.A.R. Pub., New Delhi 600-603.
- Pramar D. and Schmidt, E. L. (1989), Experimental Soil Microbiology, Burgess Publ. Co. Minnesota (USA). 15-84.
- Sable, S., P.Y. Sontakey, R. Tagade, R. D. Deotale and P. Manapure (1998), Effect of Rhizobium and molybdenum on quality aspects, yield and yield contributing characters of soybean. *Journal of Soils and Crops* **8**(2): 157-159.
- Subrahmaniyan, K., Kalaiselvan, P., Manickam, G. and Arulmozhi, N. (2000), Spacing and fertilizer requirement for confectionery groundnut varieties. *Crop Research* **19**(2): 210-12
- Subrahmaniyan, K.N., Arulmozhi, N. and Kalaiselvan, P. (1998), Effect of irrigation layout, irrigation and fertilizer level on the yield of rainfed groundnut. *Crop Research* **18**: 19-21.