

## STRONG ARCS AND MAXIMUM SPANNING TREES IN A FUZZY GRAPH

*Sameena. K and M.S. Sunitha*

### Abstract

In this paper we show that an arc is strong iff it is in at least one maximum spanning tree of a fuzzy graph.

### 1. INTRODUCTION

This paper is a continuation of the study of the concept of strong arcs in a fuzzy graph[4].

A fuzzy graph [7] is a pair  $G : (\sigma, \mu)$  where  $\sigma$  is a fuzzy subset of a set  $S$  and  $\mu$  is a fuzzy relation on  $\sigma$  such that  $\mu(u, v) \leq \sigma(u) \wedge \sigma(v)$ . We assume that  $S$  is finite and nonempty,  $\mu$  is reflexive and symmetric [7]. Also, we denote the underlying crisp graph by  $G^* : (\sigma^*, \mu^*)$  where  $\sigma^* = \{u \in S : \sigma(u) > 0\}$  and  $\mu^* = \{(u, v) \in S \times S : \mu(u, v) > 0\}$ . A fuzzy graph  $H : (\tau, \nu)$  is called a partial fuzzy subgraph of  $G : (\sigma, \mu)$ , if  $\tau \subseteq \sigma$  and  $\nu \subseteq \mu$  [6]. In particular, we call  $H : (\tau, \nu)$ , a fuzzy subgraph of  $G : (\sigma, \mu)$ , if  $\tau(u) = \sigma(u)$  for every  $u \in \tau^*$  and  $\nu(u, v) = \mu(u, v)$  for every  $(u, v) \in \nu^*$ . A path  $P$  of length  $n$  is a sequence of distinct nodes  $u_0, u_1, \dots, u_n$  such that  $\mu(u_{i-1}, u_i) > 0, i = 1, 2, \dots, n$  and the degree of membership of a weakest arc is defined as its strength. If  $u_0 = u_n$  and  $n \geq 3$ , then  $P$  is called a cycle and  $P$  is called a fuzzy cycle if it contains more than one weakest arc [5].  $G : (\sigma, \mu)$  is called a complete fuzzy graph if  $\mu(u, v) = \sigma(u) \wedge \sigma(v)$  for all  $u, v$ . The strength of connectedness between two nodes  $x$  and  $y$  is defined as the maximum of the strengths of all paths between  $x$  and  $y$  and is denoted by  $CONN_G(x, y)$ . An  $x - y$  path  $P$  is called a strongest  $x - y$  path if its strength equals  $CONN_G(x, y)$  [7].

A fuzzy graph  $G : (\sigma, \mu)$  is connected if for every  $x, y$  in  $\sigma^*$ ,  $CONN_G(x, y) > 0$ . An arc of a fuzzy graph is called strong if its weight is at least as great as the connectedness of its end nodes when it is deleted and an  $x - y$  path  $P$  is called a

---

**AMS 2000 Subject Classification:** 04A72, 05cxx

**Key words:** Strong arc, Fuzzy cycle, Complete fuzzy graph. Fuzzy tree, Maximum spanning tree.

strong path if  $P$  contains only strong arcs [4].

An arc is called a fuzzy bridge of  $G$  if its removal reduces the strength of connectedness between some pair of nodes in  $G$ . A connected fuzzy graph  $G: (\sigma, \mu)$  is a fuzzy tree if it has a partial fuzzy spanning subgraph  $T: (\sigma, \nu)$ , which is a tree, where for all arcs  $(x, y)$  not in  $T$  there exists a path from  $x$  to  $y$  in  $T$  whose strength is more than  $\mu(x, y)$  [6]. A maximum spanning tree of a connected fuzzy graph  $G: (\sigma, \mu)$  is a fuzzy spanning subgraph  $T: (\sigma, \nu)$  such that  $T^*$  is a tree and for which  $\sum_{u \neq v} \nu(u, v)$  is maximum [1].

## 2. ARCS IN A MAXIMUM SPANNING TREE

Note that if  $G$  is a fuzzy tree, then it has a unique maximum spanning tree  $T$  [9] and all arcs in  $T$  are just the strong arcs in  $G$  [4]. The propositions 4, 5 and 6 of [4] can be obtained as corollaries to the following theorem.

**Theorem 1:** An arc in a fuzzy graph  $G$  is strong if and only if it is an arc of atleast one maximum spanning tree of  $G$ .

**Proof:** Let  $(x, y)$  be a strong arc in a fuzzy graph  $G: (\sigma, \mu)$ . Then by definition,  $\mu(x, y) > 0$  and  $\mu(x, y) \geq \text{CONN}_{G-(x,y)}(x, y)$ . Consider the following two cases.

### Case I

$$\mu(x, y) > \text{CONN}_{G-(x,y)}(x, y).$$

Then, removal of  $(x, y)$  reduces the strength of connectedness between  $x$  and  $y$ , which shows that  $(x, y)$  is a fuzzy bridge in  $G$ . Then  $(x, y)$  is an arc of every maximum spanning tree of  $G$  [8].

### Case II

$$\mu(x, y) = \text{CONN}_{G-(x,y)}(x, y).$$

Then,  $(x, y)$  is an arc of some cycle  $C$  in  $G$  which contains an  $x - y$  path, say  $P$ , of strength  $\mu(x, y)$ . This implies that  $C$  contains an arc  $(u, v)$  with  $\mu(u, v) = \mu(x, y)$  and  $(x, y)$  and  $(u, v)$  are weakest arcs of  $C$ . Then it follows that there are atleast two maximum spanning trees, say,  $T_1$  and  $T_2$  with the property that  $(x, y)$  is in  $T_1$ ,  $(u, v)$  is not in  $T_1$  and  $(x, y)$  is not in  $T_2$ ,  $(u, v)$  is in  $T_2$ , which completes the proof.

Conversely, assume that  $(x, y)$  is an arc of a maximum spanning tree  $T: (\sigma, \nu)$  of  $G$  and that  $(x, y)$  is not strong. Then by definition,  $\mu(x, y) < \text{CONN}_{G-(x,y)}(x, y)$ .

This implies that there exists an  $x - y$  path, say  $P$ , whose arcs having strength greater than that of  $(x, y)$ . Then replacing the arc  $(x, y)$  by the path  $P$  in  $T$  results in another spanning tree whose total weight exceeds that of  $T$ , contradicting our assumption that  $T$  is a maximum spanning tree of  $G$ . Thus  $(x, y)$  should be strong.

Note that it also follows that all arcs in a maximum spanning tree are strong.

**Corollary 1:** If  $G$  is a fuzzy tree, then an arc of  $G$  is strong if and only if it is an arc of the maximum spanning tree of  $G$ .

**Corollary 2:**  $G$  is a fuzzy tree if and only if there is a unique strong path in  $G$  between any two nodes of  $G$ .

**Corollary 3:** In a fuzzy tree, a strong path between any two nodes  $u, v$  is a strongest  $u-v$  path.

**Remark:** Note that all arcs in a fuzzy cycle [2] and a complete fuzzy graph [3] are strong, but a fuzzy graph with all its arcs strong and whose underlying graph is complete, need not be a complete fuzzy graph as in the following example.

**Example:** Consider the fuzzy graph  $G:(\sigma, \mu)$  with  $\sigma^* = \{u, v, w, x\}$  and  $\sigma(u) = \sigma(w) = 1, \sigma(v) = .8, \sigma(x) = .5, \mu(u, v) = \mu(v, w) = .8, \mu(u, x) = \mu(x, w) = \mu(x, v) = .4, \mu(u, w) = 1$ . It can be easily verified that all arcs are strong. But  $G$  is not a complete fuzzy graph.

## REFERENCES

- [1] Bhattacharya, P. Suraweera, F. An Algorithm to Compute the Max-Min Powers and a Property of Fuzzy Graphs. *Pattern Recognition Lett.* **12** (1991) 413-420.
- [2] Bhutani, K.R., Rosenfeld, A., Fuzzy end nodes in fuzzy graphs, *Information Sciences* **152** (2003) 323-326.
- [3] Bhutani, K.R., Rosenfeld, A., Geodesics in Fuzzy Graphs, *Electronic Notes in Discrete Mathematics* **15** (2003) 51-54.
- [4] Bhutani, K.R., Rosenfeld, A., Strong arcs in Fuzzy Graphs, *Information Sciences* **152** (2003) 319-322.
- [5] Mordeson, J.N., Nair, P.S., Cycles and Cocycles of Fuzzy Graphs, *Information Sciences* **90** (1996) 39-49.
- [6] Mordeson, J.N., Nair, P.S., *Fuzzy Graphs and Fuzzy Hypergraphs*, Physica-Verlag (2000).
- [7] Rosenfeld, A., Fuzzy graphs, In *Fuzzy sets and their Applications to Cognitive and Decision Processes*, (Zadeh. L.A., Fu, K.S., Shimura, M., Eds:), Academic Press, New York (1975) 77-95.

- [8] Sunitha, M.S., Vijayakumar, A., Blocks in Fuzzy Graphs, *The Journal of Fuzzy Mathematics*, Vol. 13. No. 1 (2005) 13–23.
- [9] Sunitha, M.S., Vijayakumar, A., A characterization of Fuzzy Trees, *Information Sciences* **113** (1999) 293-300.

**Sameena K & M.S. Sunitha**

Department of Mathematics  
National Institute of Technology  
Calicut-673 601, INDIA  
([sunitha@nitc.ac.in](mailto:sunitha@nitc.ac.in))  
([y3df02@nitc.ac.in](mailto:y3df02@nitc.ac.in))