

INTO THE SEMANTICS OF QUALITY MEASURES CONCERNING FUZZY ASSOCIATION RULES

Thanuja T.S. and Mary George***

Abstract

Association rules are initially discovered in the market basket analysis [1] to identify frequently purchased items by customers. It gives certain regularities and dependencies within a data by finding frequent co-occurrence of items with a set of transactions. Usually support and confidence are the two important quality measures used to assess the quality of association rules. In fuzzy association rule mining, fuzzy support and confidence are used, which are based on t-norm operators. In addition to this opposition measures are also defined using s-norm operators. These measures were aroused during the categorization of transactions into positive and negative examples [5, 6], which we here redefined as true positive and true negative examples. Also we tried to extend these quality measures and analyzed its semantics and studied their properties.

Keywords: *Fuzzy association rules, true positive and true negative examples, Support, Opposition, fuzzy average support, fuzzy average opposition.*

1. INTRODUCTION

Data mining is the process of extracting previously unknown and potentially useful hidden predictive information from large amounts of data [1]. Association rules are initially discovered in the market basket analysis to identify frequently purchased items by customers. It give certain regularities and dependencies within a data by finding frequent co-occurrence of items with a set of transactions and relationships hidden in large data sets. The uncovered relationships can be expressed as association rules or frequent itemsets. In classical association rules, it is not possible to use every data for mining. In most real life applications, the database contains many attributes which are difficult to

* Research Scholar, Dept of Mathematics, Mar Ivanios College, Trivandrum. *Email: thanuja963@gmail.com*

** Associate Professor and Head, Dept of Mathematics, Mar Ivanios College, Trivandrum. *Email: marygeo@rediffmail.com*

represent using binary values. In such cases fuzzy sets play a major role. So in the process of association rule mining, fuzzy sets can handle both quantitative and categorical data, providing the necessary support to use uncertain data types with existing algorithms. The approach of quantitative mining allow attributes to be either members or non-members of an interval which tends to make an under or over estimation of values leading to sharp boundary problems. The use of fuzzy sets in association analysis widens the type of relationships between attributes by allowing the intervals to overlap, giving partial memberships to different sets thus avoiding unnatural boundaries in the partitioning of the attribute domain and thus making the interpretation of rules in linguistic terms easier. Thus the obtained results using fuzzy approaches are easy to understand and to apply.

An association rule is of the form $A \rightarrow B$, where A and B are attributes or sets of attributes, which tells the idea that *when A occurs in a transaction, B is likely to occur as well*. The strength of association rules can be realized by a number of quality measures. Support and confidence are the two important quality measures used essentially. Support measures the validity of an association rule where as confidence measures the quality of the rule. Thus mining association rules means, to generate all association rules $A \rightarrow B$ that have support and confidence greater than the user specified thresholds. These measures can be generalized for fuzzy association rules as well. Here we study about the transaction types and redefined their terminologies as true positive, true negative, false positive, false negative examples to understand the true semantics of the transactions. In this paper we tried to extend some of these measures, analyzed its semantics and studied their properties.

The next section explains the definition of association rules, their support and confidence measures. Section 3 gives the definition of fuzzy set and fuzzy set operations. Section 4 is devoted to fuzzy association rules, fuzzy support and confidence measures, and fuzzy average support measures. Section 5 explains the semantics of the defined measures. Section 6 contains some of the properties of the existing and defined measures and section 7 is the conclusion.

2. ASSOCIATION RULES

An association rule gives an efficient way to identify and explore certain dependencies and regularities in a database. Association rule mining was introduced by Agrawal et al. [1] as a way to discover frequently purchased items by customers in the market basket analysis.

Association rule mining is traditionally performed on a non-empty data table D with binary attributes. Let $I = \{i_1, i_2, \dots, i_n\}$ be the universe of items. Let $T = \{t_1, t_2, \dots, t_n\}$ be the transaction database (sets of objects) and let t_i represent the i^{th} transaction set in D such that $t_i \subseteq I$. Each transaction t_i is represented as a

binary vector with $t_i(i_j) = 1$ if t_i bought the item i_j and $t_i(i_j) = 0$ otherwise. An association rule is a direct association $A \rightarrow B$ with $A, B \subseteq I$ and $A \cap B = \emptyset$, where A is the antecedent and B is the consequent of the rule. A set of items X in I is called an itemset. If X contains k items, then X is called the k -itemset, T_X is the set of transactions that contain the itemset X , $|T|$ is the total number of transactions. Also we denote \widetilde{T}_A the set of transactions that does not contain the item A .

The validity and interestingness of an association rule is determined by the quality measures such as support and confidence. Support measures the extend of the simultaneous occurrence (proportion) of the items A and B in the database whereas confidence indicates the proportion of correct application of the rule.

2.1. Support Measures

Definition 2.1: The support count and respectively support of an association rule $A \rightarrow B$ is defined as:

$$supp\#(A \rightarrow B) = |T_A \cap T_B|$$

and respectively

$$supp(A \rightarrow B) = \frac{|T_A \cap T_B|}{|T|}$$

This definition of support count positive examples as it represents the transactions that explicitly support the association expressed by the rule. De Cock et al.[5, 6] classified transactions with respect of an association rule as positive example, non-positive example, negative example, non-negative example.

In order to explore the true semantics of the transaction classification, we introduce some new terminologies encouraging from the definition of confusion matrix. Thus we define

Definition 2.2: Let $A \rightarrow B$ be an association rule and t be a transaction. Then

- t is a true positive example iff $t \in T_A \wedge t \in T_B$.
- t is a true negative example iff $t \notin T_A \vee t \notin T_B$.
- t is a false positive example iff $t \in T_A \wedge t \notin T_B$.
- t is a false negative example iff $t \notin T_A \vee t \in T_B$.

This indicates how effective is our expectations. In true positive and true negative example, we got what we expect, according as presence or absence of items. In false positive examples we assume the presence of some items, but it was a false one and in false negative examples, we assume the absence of some items and it appeared to be false.

Based on this classification, we get the following different measures:

Definition 2.3: Let $A \rightarrow B$ be an association rule. Then

1. minimum support count: $minsupp\#(A \rightarrow B) = |T_A \cap T_B|$
2. maximum opposition count: $maxopp\#(A \rightarrow B) = |\widetilde{T}_A \cup \widetilde{T}_B|$
3. minimum opposition count: $minopp\#(A \rightarrow B) = |T_A \cap \widetilde{T}_B|$
4. maximum support count: $maxsupp\#(A \rightarrow B) = |\widetilde{T}_A \cup T_B|$

and the corresponding measures is given by

1. minimum support : $minsupp(A \rightarrow B) = \frac{|T_A \cap T_B|}{|T|}$
2. maximum opposition count: $maxopp\#(A \rightarrow B) = \frac{|\widetilde{T}_A \cup \widetilde{T}_B|}{|T|}$
3. minimum opposition count: $minopp\#(A \rightarrow B) = \frac{|T_A \cap \widetilde{T}_B|}{|T|}$
4. maximum support count: $maxsupp\#(A \rightarrow B) = \frac{|\widetilde{T}_A \cup T_B|}{|T|}$

Remark 2.4

1. $minsupp(A \rightarrow B) \leq maxsupp(A \rightarrow B)$
2. $minopp(A \rightarrow B) \leq maxopp(A \rightarrow B)$

2.2. Confidence Measures

Definition 2.5: The confidence of a rule $A \rightarrow B$ is defined as:

$$conf(A \rightarrow B) = \frac{supp\#(A \rightarrow B)}{supp\#(A)}$$

Confidence can be treated as the conditional probability ($P(B|A)$) or the relative cardinality of B with respect to A .

Definition 2.6: (Hullermeier, [9]) The confidence measure, n – confidence is defined as:

$$conf_n(A \rightarrow B) = \frac{minsupp\#(A \rightarrow B)}{minopp\#(A \rightarrow B)}$$

Definition 2.7: (DeCook et al.,[5]) The pessimistic confidence p – confidence and the optimistic confidence o – confidence are defined as:

$$conf_p(A \rightarrow B) = \frac{minsupp\#(A \rightarrow B)}{maxopp\#(A \rightarrow B)}$$

$$conf_o(A \rightarrow B) = \frac{maxsupp\#(A \rightarrow B)}{minopp\#(A \rightarrow B)}$$

Definition 2.8: ([11]) Given a pair (M_1, M_2) of quality measures for association rules, with the property $M_1(A \rightarrow B) \leq M_2(A \rightarrow B)$, the inferior confidence and superior confidence are defined as

(a) inferior confidence

$$conf_*(A \rightarrow B) = \frac{\alpha \cdot M_1(A \rightarrow B)}{(1 - \beta) \cdot M_1(A \rightarrow B) + \beta \cdot M_2(A \rightarrow \tilde{B})}$$

(b) superior confidence

$$conf^*(A \rightarrow B) = \frac{\alpha \cdot M_2(A \rightarrow B)}{(1 - \beta) \cdot M_1(A \rightarrow B) + \beta \cdot M_1(A \rightarrow \tilde{B})}$$

Remark 2.9

$$conf_p(A \rightarrow B) \leq conf_n(A \rightarrow B) \leq conf_o(A \rightarrow B) \\\ conf_*(A \rightarrow B) \leq conf^*(A \rightarrow B)$$

3. FUZZY SETS AND FUZZY SET OPERATIONS

A fuzzy set A in a given universal set X is a mapping from $X \rightarrow [0,1]$, usually denoted as $A = \{ (x, A(x)): x \in X \}$ where $A(x)$ is called the grade of membership of each $x \in X$. The cardinality of a fuzzy set A in X is defined as $|A| = \sum_{x \in X} A(x)$.

A monotonic, associative and commutative mapping from $[0,1]^2 \rightarrow [0,1]$ is called t -norm T , if it satisfies $T(x, 1) = x$ for all $x \in [0,1]$ and a t -conorm S if it satisfies $S(x, 0) = x$ for all $x \in [0,1]$. A fuzzy complement N is a decreasing mapping from $[0,1] \rightarrow [0,1]$ satisfying $N(0) = 1$ and $N(1) = 0$.

For the fuzzy sets A and B in X , the complement, intersection and union can be defined by

$$co A(x) = \tilde{A}(x) = N(A(x)) \\ (A \cap_T B)(x) = T(A(x), B(x)) \\ (A \cup_S B)(x) = S(A(x), B(x))$$

4. FUZZY ASSOCIATION RULES

For quantitative and categorical attributes, binary rules are not satisfactory. As association rule mining deals with only binary transaction data, a new approach emerged out using fuzzy sets to mine quantitative data frequently present in databases efficiently, called fuzzy association rules.

Let $T = \{t_1, t_2, \dots, t_n\}$ be the transaction database (sets of objects) and let t_i represents the i^{th} transaction in D . Let $I = \{i_1, i_2, \dots, i_m\}$ be the universe of items. Each attribute i_k will associate with several fuzzy sets. In order to represent the fuzzy sets associated with i_k , we use the notion $F_{i_k} = \{f_{i_k}^1, f_{i_k}^2, \dots, f_{i_k}^l\}$ where $f_{i_k}^j$ is the j^{th} fuzzy set in F_{i_k} .

Definition 4.1: A fuzzy association rule is of the form: $(X \in F_X) \rightarrow (Y \in F_Y)$ where $X, Y \subset I, X \cap Y = \emptyset, X = \{x_1, x_2, \dots, x_p\}$ and $Y = \{y_1, y_2, \dots, y_q\}$ are attributes, and $F_X = \{f_{x_1}, f_{x_2}, \dots, f_{x_p}\}$ and $F_Y = \{f_{y_1}, f_{y_2}, \dots, f_{y_q}\}$ are fuzzy sets that characterize X and Y respectively. For each fuzzy set $f_{i_k}^l$ we can associate a membership function $\mu(f_{i_k}^l): dom(i_k) \rightarrow [0,1]$ corresponding to the attribute i_k . Fuzzy sets and their corresponding membership functions have to be defined by domain experts.

Now we define the support and confidence measures of fuzzy association rules, extending the crisp association rules. Here we work on simple association rules $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ in which A and B are both attributes, not the sets of attributes.

4.1. Fuzzy Support Measures

Definition: 4.2: The fuzzy support count and respectively fuzzy support of a fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is usually defined as:

$$f_{supp\#}(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \sum_{x \in T} F_A \cap_T F_B(x)$$

and respectively

$$f_{supp}(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{\sum_{x \in T} F_A \cap_T F_B(x)}{|T|}$$

Definition 4.3: Let $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ be a fuzzy association rule. Then we define:

(a) fuzzy minimum support:

$$f_{minsupp}(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{\sum_{x \in T} (F_A \cap_T F_B)(x)}{|T|}$$

(b) fuzzy maximum opposition:

$$f_{maxopp}(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{\sum_{x \in T} (\widetilde{F_A} \cup_S F_B)(x)}{|T|}$$

(c) fuzzy minimum opposition:

$$fminopp(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{\sum_{x \in T} (F_A \cap_T \widetilde{F_B})(x)}{|T|}$$

(d) fuzzy maximum support:

$$fmaxsupp(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{\sum_{x \in T} (\widetilde{F_A} \cup_S F_B)(x)}{|T|}$$

4.2. Fuzzy Average Support Measures

Studying the above definitions in view of averaging operators we define another two measures, fuzzy average support and fuzzy average opposition of the fuzzy association rule.

Definition 4.4: The fuzzy average support and fuzzy average opposition of a fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is defined as:

(a) Fuzzy average support:

$$\begin{aligned} favrgsupp(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) \\ = \frac{\sum_{x \in T} [\lambda(\widetilde{F_A} \cup_S F_B)(x) + (1 - \lambda)(F_A \cap_T F_B)(x)]}{|T|} \end{aligned}$$

where, $\lambda \in [0,1]$.

(b) fuzzy average opposition:

$$\begin{aligned} favrgopp(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) \\ = \frac{\sum_{x \in T} [\lambda(F_A \cup_S \widetilde{F_B})(x) + (1 - \lambda)(F_A \cap_T F_B)(x)]}{|T|} \end{aligned}$$

where, $\lambda \in [0,1]$.

Similarly we can define the corresponding count measures: $fminsupp\#, fmaxopp\#, fminopp\#, fmaxsupp\#, favrgsupp\#, favrgopp\#$.

4.3. Fuzzy Confidence Measures

Definition 4.5: The fuzzy confidence of a fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is defined as

$$fconf(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{\sum_{x \in T} (F_A \cap_T F_B)(x)}{\sum_{x \in T} F_A(x)}$$

Now we define fuzzy version of $conf_n, conf_p, conf_o$ defined by Hullermeier, [9] and DeCook et al.,[5].

Definition 4.6: The fuzzy confidence measures n – confidence, p – confidence and o – confidence of a fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is defined as:

(a) fuzzy n – confidence

$$fconf_n(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{fminsupp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}{fminopp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}$$

(b) fuzzy pessimistic confidence

$$fconf_p(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{fminsupp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}{fmaxopp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}$$

(c) fuzzy optimistic confidence

$$fconf_o(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{fmaxsupp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}{fminopp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}$$

Now we define two more confidence measure fuzzy m – confidence and fuzzy mn – confidence

Definition 4.7: The fuzzy confidence measure m – confidence of a fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is defined as:

$$fconf_m(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{fmaxsupp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}{fmaxopp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}$$

Definition 4.8: The fuzzy confidence measure mn – confidence of a fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is defined as:

$$fconf_{mn}(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) = \frac{favrsgsupp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}{favrgopp\#\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle}$$

Remark 4.9:

(i) For $\lambda = 0$, $fconf_{mn} = fconf_n$

(ii) For $\lambda = 1$, $fconf_{mn} = fconf_m$

Definition 4.10: [Refer Definition 2.8] The fuzzy inferior confidence and respectively fuzzy superior confidence of the fuzzy association rule $\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$ is defined as:

(a) fuzzy inferior confidence

$$fconf_*\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle = \frac{\alpha . M_1(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle)}{(1 - \beta) . M_1(\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) + \beta . M_2(\langle A, F_A \rangle \rightarrow \langle B, \widetilde{F}_B \rangle)}$$

(b) superior confidence

$$fconf^* \langle A, F_A \rangle \rightarrow \langle B, F_B \rangle$$

$$= \frac{\alpha \cdot M_2 (\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle)}{(1 - \beta) \cdot M_2 (\langle A, F_A \rangle \rightarrow \langle B, F_B \rangle) + \beta \cdot M_1 (\langle A, F_A \rangle \rightarrow \langle B, \widetilde{F}_B \rangle)}$$

with $\alpha, \beta \in [0,1]$.

5. INTO THE SEMANTICS OF THE DEFINED MEASURES

Based on the confusion matrix terminology Table1, we divide the transaction into true positive (TP), true negatives (TN), False Positive (FP) and False Negative (FN) examples. For an association rule $\rightarrow B$, where A is the antecedent and B is the consequent of the rule, TP is the number of instances which match with rule antecedent and consequent, TN is the number of instances which match rule antecedent and consequent FP is the number of instances which match only with rule antecedent and FN is the number of instances which match only with rule consequent.

Table 1: Confusion Matrix

Actual class	Predicted class	
	YES	NO
YES	TP: True Positive	FP: False Negative
NO	FP: False Positive	TN: True Negative

So we can view $fminsupp$ as the rate of true positive example, $fmaxopp$ as the rate of true negative example, $fminopp$ as the rate of false positive example, $fmaxsupp$ as the rate of false negative example. Thus $fminopp$ is termed as positive error rate and $fmaxsupp$ is termed as negative error rate. $favrgsupp$ and $favrgopp$ represent the rate of presence and absence of items respectively. Also using different fuzzy confidence measures which are defined using fuzzy support measures, we can effectively explains the precision, specificity, sensitivity of each association rules.

6. PROPERTIES

6.1. Properties of Fuzzy Support and Confidence Measures

Remark 6.1: The measures $fconf_*$ and $fconf^*$ defined above represent a general and integrated frame for the quality measures as most of them can be obtained particularizing $T, S, N, \alpha, \beta, M_1, M_2$. Thus we get the following

- (a) For $M_1 = M_2 = fsupp\#, \alpha = \beta = \frac{1}{2}, T(a, b) = ab$ and $N(a) = 1 - a$, we have:

$$fconf_* = fconf^* = fconf$$

- (b) For $M_1 = M_2 = fsupp\#, \alpha = \beta = 1$ we have:

$$fconf_* = fconf^* = fconf_n$$

- (c) For $M_1 = fminsupp\#, M_2 = fmaxsupp\#$ and $\alpha = \beta = 1$ we have:

$$fconf_* = fconf_p$$

$$fconf^* = fconf_o$$

- (d) For $M_1 = fminsupp\#, M_2 = fmaxsupp\#$ and $\alpha = \beta = \frac{1}{2}$ we have:

$$fconf_* = fminsupp$$

$$fconf^* = fmaxsupp$$

- (e) For $M_1 = fminopp\#, M_2 = fmaxopp\#, \alpha = \beta = \frac{1}{2}$ and $N(a) = 1 - a$ we have:

$$fconf_* = fminopp$$

$$fconf^* = fmaxopp$$

6.2. Properties of Fuzzy Average Support and Fuzzy Average Opposition Measures

Using the new measures, fuzzy average support and fuzzy average opposition we arrive at the following:

- (a) For $M_1 = M_2 = favrgsupp\#, \alpha = \beta = \frac{1}{2}, T(a, b) = ab, S(a, b) = a + b - ab$ and $N(a) = 1 - a$, we have:

(i) For $\lambda = 0, fconf_* = fconf^* = fconf$

(ii) For $\lambda = 1,$

$$\frac{1}{fconf_*} = \frac{1 - fminsupp}{1 - fminopp} + 1 = \frac{1}{fconf^*}$$

- (b) For $M_1 = M_2 = favrgsupp\#$ and $\alpha = \beta = 1,$ we have

(i) For $\lambda = 0, fconf_* = fconf^* = fconf_n = fconf_{mn}$

(ii) For $\lambda = 1, fconf_* = fconf^* = fconf_m = fconf_{mn}$

(c) For $M_1 = M_2 = favrgopp\#, \alpha = \beta = \frac{1}{2}, T(a, b) = ab, S(a, b) = a + b - ab$ and $N(a) = 1 - a$, we have:

(i) For $\lambda = 0, fconf_* = fconf^* = 1 + fconf$

(ii) For $\lambda = 1,$

$$\frac{1}{fconf_*} = \frac{1 - fminopp}{1 - fminsupp} + 1 = \frac{1}{fconf^*}$$

(d) For $M_1 = M_2 = favrgopp\#$ and $\alpha = \beta = 1,$ we have

(i) For $\lambda = 0,$

$$fconf_* = fconf^* = \frac{1}{fconf_n} = \frac{1}{fconf_{mn}}$$

(ii) For $\lambda = 1,$

$$fconf_* = fconf^* = \frac{1}{fconf_m} = \frac{1}{fconf_{mn}}$$

7. CONCLUSION

In this paper we extend some quality measures defined for crisp association rule to fuzzy association rule. Here we study about the transaction types and redefined their terminologies as true positive, true negative, false positive, false negative examples to understand the true semantics of the transactions. Based on this, we defined various support and confidence measures and extended to fuzzy average support and opposition measures. We also analyzed the semantics of these measures and studied some of the properties of the newly defined measures comparing the existing ones.

Acknowledgment

This research was supported by the Kerala State Council for Science, Technology and Environment (KSCSTE).

References

- [1] R. Agrawal, T. Imielinski, and A. Swami Mining Association Rules between Sets of Items in Large Databases, In Proceedings of the International Conference on Management of Data, Washington, D.C., pp. 207-216, May 1993.
- [2] R. Agrawal, R. Srikant, Fast Algorithms for Mining Association Rules, In Proc. 20th Int. Conf. Very Large Data Bases, VLD, Santiago, Chile, 1994, pp.487-499.
- [3] Bakk. Likas Helm, Fuzzy Association Rules: an implementation in R, Vienna University of Economics and Buiseness Administration, 2007, Vienna.

- [4] Chan, Keith C.C.; Au, Wai-Ho, An Effective Algorithm for Discovering Fuzzy Rules in Relational Databases, In Proceedings of Fuzzy Systems, IEEE World Congress on Computational Intelligence, 1998.
- [5] M. De Cock, C. Cornelis, E. E. Kerre, Fuzzy Association Rules: a Two-Sided Approach, In: Y. Liu, G. Chen and K. Y. Cai (Eds.): Proceedings of FIP2003 (International Conference on Fuzzy Information processing: Theories and Applications), Tsinghua Univ. Press, 2003, pp. 385-390.
- [6] M. De Cock, C. Cornelis, E.E. Kerre, A clear view on quality measures for fuzzy association rules, In Proceedings of Int. Conf. on Fuzzy Sets and Soft Computing in Economics and Finance, St. Petersburg, 2004, pp. 54-61.
- [7] D. Dubois, E. Hüllermeier and H. Prade, A Note on Quality Measures for Fuzzy Association Rules, In Proceedings IFSA-03, 10th International Fuzzy Systems Association World Congress, Lecture Notes in Artificial Intelligence, number 2715, Springer-Verlag, 2003, pp. 346-353.
- [8] George J.Klir and Bo Yuan, Fuzzy Sets and Fuzzy Logic: Theory and Applications, Pearson Education, 1995, USA.
- [9] E. Hüllermeier, Fuzzy Association Rules: Semantic Issues and Quality Measures, In: B. Reusch (Ed.) Proceedings of the International Conference, 7th Fuzzy Days on Computational Intelligence, Theory and Applications, Lecture notes in Computer Science, number 2606, 380-391, Springer-Verlag.
- [10] Hong T P, Lin K Y, Chien B C, Mining fuzzy multiple-level association rules from quantitative data, Applied Intelligence, 18(1): 79-90,2003.
- [11] I.Iancu, M. Gabroveanu, A. Giurca, A Pair of Confidence Measures for Association Rules, In: 30th Annual Conference of the German Classification Society, GfKI 2006, Berlin, March 8-10, 2006 (in press).
- [12] Kuok C, Fu A, Wong M, Mining fuzzy association rules in databases, SIGMOD Record, 27(1):41-46, 1998.
- [13] Li-Xin Wang, A Course in Fuzzy Systems and Control, Prentice Hall International, Inc., 1997, Hong Kong.