

Studies on Effect of Blanching on Turmeric Processing

¹R. R. Patil., ²S. P. Kurhekar, ³S. R. Patil and ⁴S. R. Sawant

ABSTRACT: Turmeric is valued for its deep yellow colour and aromatic flavour. Processing of turmeric assumes importance from of appearance and colour point of view. The processing of turmeric by blanching was compared to the conventional method of boiling the rhizomes. It was observed that the average length, breadth and thickness of fresh turmeric rhizomes was 71.29, 25.19 and 24.08mm whereas that of blanched turmeric rhizomes it was 69.72, 24.42 and 22.99mm respectively. The average values of weight, volume, and true density of fresh and blanched single turmeric rhizome were observed to be 26.86g, 26.15cm³, 1.026 g/cm³ and 25.62g, 24.23cm³, 1.07g/cm³, respectively. The skin removal and stick piercing in turmeric rhizomes blanched for 25, 30 and 35 minutes is less easy, easy and more easy respectively. The turmeric rhizomes blanched and boiled for 25, 30 and 35 minute retains curcumin content 4.27%, 4.09%, 4.01% and 4.21%, 3.91%, 2.29% respectively

Keywords: turmeric, blanching

INTRODUCTION

Turmeric is an important cash crop. It has a good demand in India and other countries. It is a herbaceous, perennial crop with thick underground rhizomes giving rise to primary and secondary rhizomes called fingers. Turmeric is valued for its deep yellow colour and aromatic flavour. It is used both as a spice and as a food colouring agent in pickles, chutneys, curries and other culinary preparations. It is also used as a drug and cosmetic. Processing of turmeric assumes importance from of appearance and colour point of view. The core of turmeric is reddish which after boiling becomes, yellow and of uniform in colour. Curing is the process of cooking raw rhizomes in hot water to obtain attractive colour, characteristics aroma, destroy and viability of fresh rhizomes and obviate the raw odour, reduces the time of drying, ensures an even distribution of colour in the rhizomes and gives better quality product by gelatinization of starch In steam blanching, water vapour hardly escapes in the atmosphere, steam is uniformly distributed throughout the mass and uniform blanching is facilitated. Loss of heat is prevented, easy handling and cost of processing is

reduced. The overall cost of processing will be reduced due to above reasons and due to faster rate of blanching quality of turmeric will also be improved.

MATERIALS AND METHOD

1. Turmeric Steam Blancher.
2. Raw Turmeric (Variety Salem)
3. Gunny bags (for transporting and storing raw turmeric)
4. Drying Yard
5. The apparatus used are glass wares, test tube, sieve, Volumetric Flask, Chemical such as alcohol, acetone etc.
6. Wood was used as fuel.

Determination of Physical Properties

(a) *Size and Shape:* The maximum tri-axial dimensions i.e. length, breadth and thickness were obtained from 20 randomly selected turmeric rhizomes using a vernier caliper. The samples were divided into two categories i.e., fresh / raw and cure at atmospheric pressure. The averages of the readings were taken as its length, breadth and thickness.

1. Assistant Professor, Department of Farm Structures, College of Agricultural Engineering and Technology, DBSKKV, Dapoli.
2. Ex Faculty Member, Sau KSK College of Food Technology, Beed.
3. Ph.D Scholar, Karunya University, Karunya Nagar.
4. B.Tech Student, College of Agricultural Engineering and Technology, DBSKKV, Dapoli.

(b) **Volume and True density:** The volume and true density of fresh /raw turmeric rhizomes as well as cured rhizomes were measured by toluene displacement method.

Blanching Process

Soil and other particles adhering to rhizomes were removed by hand cleaning and washing before blanching under clean running water. Roots and fibers were trimmed off. After that the mother and finger rhizomes were separated and weighted. Then the rhizomes were loaded manually into the blanching vessels and the lid at the top of vessels were closed and tightened by nut and bolt arrangement.

The water tank was filled with clean water from water inlet situated at the top of water tank. Fuel was fed to the furnace. Furnace provides space for combustion of fuel. The combustion of fuel heats up the water in the water tank. The flue gases passes through the fire tubes surrounded by water in the tank up to the chimney. After reaching the water temperature to 100°C, steam was generated which rises up and gets collected in the steam chamber.

Steam conduit conveys the steam from the steam chamber to the blanching vessel. After reaching the steam pressure to 0.1kg/cm² in the boiler the valve of first vessel was opened and steam was diverted into the blanching vessel. In the blanching, the steam was distributed by four long vertical perforated pipes. The rhizomes were blanched for 25, 30 and 35 min. with the help of timer. The pressure and temperature were regulated duration this period with the help of control valve and fuel feed rate. Pressure gauge indicates the pressure in the vessels. The blanching was carried out batch wise in the vessel alternatively. The rhizomes in one vessel were blanched and during this interval the rhizomes in other vessel were unloaded in the trolley by opening the shutter with the help of wheel located at the bottom of vessel. The trolley full of blanched turmeric can be pushed by one person to drying yard to spread. Blanched turmeric was spread on the drying yard near by end and allowed to dry upto 6% moisture content. The procedure was repeated for different time interval and different observation were noted and tabulated.

Turmeric Boiling

For comparative study the 3 quintal turmeric rhizomes were boiled for 25,30, and 35 minutes

Chemical Analysis

About 0.1 to 0.2 gm. finely ground turmeric powder was extracted by refluxing over a water condenser with 40 ml of distilled alcohol for 2.5 hours. The extract was transferred to 100 ml, volumetric flask and made to volume with alcohol. It was then filtered and an aliquot of 5 ml was transferred to a 100ml volumetric flask. It was mixed well and the absorbance of this solution was measured at 425 nm. against alcohol blank. Using the absorbance value of standard solution of curcumin (0.0025g/100ml gives and absorbance of 0.42) the curcumin percentage was calculated

RESULT AND DISCUSSION

Effect of Blanching on Physical Properties of Turmeric

The average length, breadth and thickness of fresh turmeric rhizomes was 71.29, 25.19 and 24.08mm whereas that of blanched turmeric rhizomes it was 69.72,24.42 and 22.99mm respectively. The average size of fresh and blanched turmeric rhizomes is 34.51 mm and 33.88 mm respectively. The sphericity of fresh and blanched turmeric rhizomes is 0.495 and 0.489. The average values of weight, volume, and true density of fresh and blanched single turmeric rhizome were observed 26.86g, 26.15cm³, 1.026 g/cm³ and 25.62g, 24.23cm³, 1.07g/cm³, respectively.

The decrease in the axial dimensions is due to oozing out of moisture in the form of vapour at the temperature 100°C in the blanching vessels. No water droplets observed on the blanched turmeric rhizomes after taking out of vessels as in the traditionally boiled turmeric in boiling pots.

Effect of Blanching Time on Turmeric Rhizomes

It was observed that the skin removal and stick piercing in turmeric rhizomes blanched for 25,30 and 35 minutes is less easy, easy and more easy

Table 1
Average length, breadth, thickness, size, sphericity and true density of fresh and cured turmeric rhizomes.

Turmeric	Length (mm) <i>a</i>	Breadth (mm) <i>b</i>	Thickness (mm) <i>c</i>	Size (mm) $(abc)^{1/3}$	Sphericity $(abc)^{1/3} a$	Weight (g)	Volume (cm ³)	True density g/cm ³
Fresh Av.	71.29	25.19	24.08	34.51	0.495	26.86	26.15	1.03
Cured Av.	69.72	24.42	22.99	33.88	0.489	25.62	24.23	1.07

Table 2
Effect of blanching time on turmeric rhizomes.

Time interval (Minutes)	Skin removal	Stick piercing	Colour Uniformity
25	Less easy	Less easy	Non uniform yellow, Core is visible
30	Easy	Easy	Uniform yellow
35	More easy	More easy	Uniform yellow, very soft rhizomes observed

respectively. It is also observed that turmeric blanched for 25 minutes attains non uniform yellow colour with separate core and outer layer . Turmeric rhizomes blanched for 30 and 35minute attains uniform yellow colour but the rhizomes becomes very soft after 35minutes. Turmeric boiled for 25 minutes was not boiled uniformly.

Effect of method of turmeric processing on curcumin and oleoresin content

It was observed that the turmeric rhizomes blanched and boiled for 25, 30 and 35 minute retains curcumin content 4.27%, 4.09%, 4.01% and 4.21%, 3.91%, 2.29% respectively. Also the values of oleoresin content in the rhizomes blanched and boiled for the above time period were observed 5.01%, 4.92%, 4.90% and 4.98%, 4.86% and 4.81% respectively. The of loss curcumin and oleoresin content in turmeric boiling is more as compared to steam blanching.

Table 3
Effect of blanching time on curcumin and Oleoresin content

Content %	Time interval in minutes					
	Steam blanching			Boiling		
	25	30	35	25	30	35
Curcumin %	4.27	4.09	4.01	4.21	3.91	2.29
Oleoresin %	5.01	4.92	4.90	4.98	4.86	4.81

Effect of blanching on drying of turmeric

The initial moisture content of turmeric was found to be 76%. After blanching the moisture content of steam

blanched and boiled turmeric was observed as 74.5% and 77% respectively. The time of drying for blanched and boiled turmeric was 8 days and 12 days respectively

CONCLUSIONS

1. The quality of turmeric steam blanched for 30 minutes is better than boiled turmeric with respect to curcumin and oleoresin percentage retained.
2. Time required for drying of blanched turmeric was (8days) less than boiled turmeric(12days).
3. Loss of fuel, labour, time and quality can be reduced by using Turmeric Steam Blancher.

REFERENCES

Krishanmurthy, M.N., Padmabai, R., Natrajan, C.P. and Kuppaswamy, S. (1995), Colour content of turmeric varieties and studies on its processing. J. Food Sci. Technol., 12:12.

Mohesnin, N.N. (1986), Physical properties of plant and Anima Materials. Gordon and Breach Science Publishers, New York.

Nair, M.K. (1984), Turmeric selection. Central Plantation Crops Research Institute (Private Communication).

Varshney, A.K., Garala, S.N. and Akbari, S.H. (2004), Effect of curing on physical characteristics of turmeric J. Agril. Engg., 2:16-19.

Wanichgarnjaakul (1996), Effect of curing on turmeric tissue and drying time. kasetart. J. Natural Sci., 30 (4): 485-492.

