

## Pathogenicity and Morphological Variabilities of *Lasiodiplodiatheobromae* Isolates in Tuberose

D. Durgadevi. A. Sankaralingam and S. Prabhu

**ABSTRACT:** Peduncle blight, hitherto an unknown disease was found to be a major limiting factor to the cultivation of tuberose, as the disease incidence was noticed up to 42.60 per cent in pockets of Madurai district. Though *Lasiodiplodia theobromae* is an ubiquitous pathogen, its occurrence on tuberose is a new record. Peduncle blight disease infected tuberose plants were collected from seven places of Tamil Nadu, India. Among the seven isolates screened for virulence, isolate I<sub>1</sub> collected from Agricultural College in Madurai district recorded maximum infection under artificial inoculation. The isolate I<sub>1</sub> significantly recorded 570 mm lesion length in pinpricked buds and also recorded 47.00 mm mycelial growth per day. *Lasiodiplodiatheobromae* produced dark brown, flask-shaped, ostiolate pycnidia, 110-170 mm x 60-130 mm in size appeared in seven-day-old cultures. Conidia were initially globose to oblong, hyaline and unicellular later turning brown and septate, measuring 18.5-21.7 mm x 8.0-11.2 mm. Compact, fluffy and sparse colony types were observed among the isolates. Significant variation was observed in morphological character and virulence among the *Lasiodiplodia theobromae* isolates.

**Key words:** Morphological character, *Lasiodiplodia theobromae*, Pathogenicity, Tuberose

### INTRODUCTION

Tuberose (*Polianthes tuberosa* Linn.) is one of the most important ornamental plants which is extensively cultivated in many sub-tropical and tropical areas of the world (Biswas *et al.*, 2002) Total area under tuberose in India is around 3000 ha with an average yield of 6300 kg per ha. The flower production in Maharashtra was 9180 tonnes (Bhattachaarjee *et al.*, 1994). In India the incidence of peduncle blight disease was first reported by (Durgadevi and Sankaralingam, 2012). The causal organism *Lasiodiplodia theobromae* was found to be associated with this disease, producing blossom blight, peduncle blight and blighting of leaf tips.

Morphological characters are important tools in the identification and classification of fungus. Several workers described enormous pathogenic variability among *Lasiodiplodia theobromae* isolates (Patil *et al.*, 2006; Khanzada *et al.*, 2006). Hence, the present investigation was conducted to study the morphological and pathogenic variability among isolates of *Lasiodiplodia theobromae*.

### ISOLATION OF PATHOGEN

Peduncle blight disease infected plants were collected from tuberose growing areas of The pathogen causing peduncle blight in tuberose was isolated from the samples by tissue segment method Infected portions of diseased plants were cut into small pieces using sterilized scalpel and surface sterilized with 0.1 per cent mercuric chloride for one minute and washed in three changes of sterile distilled water and then placed on Petri dish containing solidified Potato Dextrose Agar (PDA) medium. These plates were incubated at room temperature (28 ± 2°C) for five days and observed for the growth of the fungus. The hyphal tips of fungi were transferred aseptically to PDA slants for maintenance of the culture on potato dextrose agar (PDA) and the fungus was purified by single spore isolation and maintained on PDA. The causal organism was identified based on spore morphology and confirmed further (ID.NO. 6751/11) by Indian Type Culture Collection Centre (ITCC) of Division of Plant Pathology, Indian Agricultural Research Institute, New Delhi.

\* Department of Plant Pathology, Center for Plant protection studies, Tamil Nadu Agricultural University, Coimbatore - 641003, Tamil Nadu, India, E-mail: dev\_i.agri18@gmail.com

## MORPHOLOGICAL VARIABILITY AMONG *LASIDIPLODIA THEOBROMAE* ISOLATES

From the five day old culture plates nine mm culture disc of the pathogen was cut by a sterilized cork borer and placed at the center of each sterile Petri dish containing 20 ml of previously sterilized and solidified PDA medium. The plates were incubated at room temperature (28±2°C) for five days. The growth and morphological characters of the isolates viz., colony morphology, mycelial growth rate, colony colour, pycnidia size, shape and septation were observed, measurement was made under the microscope (Olympus BX41).

## PATHOGENIC VARIABILITY AMONG *LASIDIPLODIA THEOBROMAE* ISOLATES

### Detached Flower Bud Technique

A five- mm culture disc of *L. theobromae* was placed closer to the calyx of healthy detached flower bud and kept in 150-mm-dia Petri dish over a layer of moistened filter paper. An empty five- mm disc of PDA served as control.

Three replications were maintained and the plates were incubated at room temperature (28 ± 2° c). The formation of lesion on flower bud was closely monitored and the lesion length was recorded at regular intervals.

### Pathogenicity in Glasshouse

The pathogenicity of the fungus was confirmed by Koch's postulates using five numbers of four-month-old healthy plants. Plants were inoculated by making a vertical cut (3 mm) in the peduncle region below the calyx using a sterilized needle and placing a fungal disc over the wound. The inoculated area was covered with moist cotton and wrapped with parafilm. The plants were covered with polythene

bags to maintain humidity and monitored for symptom expression. Proper controls were maintained with PDA plugs.

## RESULTS AND DISCUSSION

The growths and colony characters of seven isolates of *L. theobromae* were assessed on PDA. Of the seven isolates studied for their cultural characters, five isolates, namely I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub>, I<sub>4</sub> and I<sub>5</sub>, showed colony growth rate of 1.63 to 1.88 mm/h indicating their fast growing nature (Table 1, Figure 1a, 1b). The isolate I<sub>2</sub> attained full nine-cm-dia growth at 48 h after inoculation. On the same day growth of the other isolates I<sub>2</sub> (88.30mm), I<sub>3</sub> (87.00mm) and I<sub>4</sub> (86.60mm) were on par with I<sub>1</sub>. The lowest growth was observed with the isolate I<sub>6</sub> (64.30 mm). The colony colour of the isolates were dull white (I<sub>1</sub>, I<sub>7</sub>), cottony white changing to black (I<sub>4</sub>, I<sub>6</sub>) and greyish white (I<sub>2</sub>, I<sub>5</sub>). The isolates I<sub>1</sub>, I<sub>4</sub> and I<sub>6</sub> showed dense, fluffy growth. I<sub>3</sub> was partial fluffy. However, I<sub>2</sub> and I<sub>4</sub> were sparse in growth. The growth of I<sub>5</sub> was appressed. The mycelial were fast spreading, branched, septate and pycnidia were brown coloured. Conidia were initially globose to oblong, hyaline and unicellular turning brown and septate later. This is in accordance with the descriptions of *L. theobromae* reported earlier by Punithalingam (1979). The results indicated that difference in morphological character was positively correlated with its virulence.

Peduncles of tuberose showing typical symptoms of blight were collected from seven locations (Table 2). The age of the crop varied from 9 to 28 months and cultivars with single flower were common than double flower types. The disease incidence ranged from 12.00 to 42.60 per cent. In the current survey, blighting of flower buds of tuberose followed by dieback of peduncle from tip downward are the major

Table 1  
Morphological Characters and growth of Different Isolates of *L. theobromae* on Potato Dextrose Agar

Sl. No.	Isolate code	Colony type	Colour	Mycelial growth	
				(mm)	rate (mm/h)
				48 h	
1	I <sub>1</sub>	Coarse, fluffy	Dull white	90.00	1.88
2	I <sub>2</sub>	Sparse	Greyish white	88.30	1.84
3	I <sub>3</sub>	Partial fluffy	Dull white	87.00	1.81
4	I <sub>4</sub>	Coarse	Black	86.60	1.80
5	I <sub>5</sub>	Flat	Greyish white	78.30	1.63
6	I <sub>6</sub>	Dense, Fluffy	Black	64.30	1.34
7	I <sub>7</sub>	Sparse	Dull white	77.00	1.60
		CD		6.44	





Figure 1a: Isolates of *Lasiodiplodia theobromae* on PDF Growth at 3 Days after inoculation



1b: View through the bottom of the culture plates

symptoms observed in the field. Tests of pathogenicity by detached flower technique (Figure 2) as well as those conducted at greenhouse yielded symptoms as observed in the field. The fungus also caused symptoms on flower buds without wound after an incubation period of seven days. However, wounding was found to enhance the symptom expression.


The fungus *Lasiodiplodia theobromae* has a wide host range infecting monocot and dicot plants causing an array of symptoms including shoot bight and die back

(Arx, 1987). In cashew it causes drying of petals and other flower parts followed by die back of peduncles leading to inflorescence blight (Olunloyo, 1979). The fungus causes necrosis and die back of shoots in mango (Burhan, 1987; Khanzada *et al.*, 2004) and grapevine (Wood and Wood, 2005).

The present investigation revealed that the isolates of *Lasiodiplodia theobromae* exhibited high variability in morphological character and pathogenicity which could be used for further development of race specific resistant varieties of onion for the control of disease.

Table 2  
Occurrence of Peduncle Blight in Tuberose Growing Areas

Sl. No.	Isolate code	Location	Variety	Age of the crop (months)	Disease Incidence (%)
1	I <sub>1</sub>	AC and RI (Madurai)	Suvasini (Double)	24	42.60
2	I <sub>2</sub>	Vilampatti (Dindugaldt)	Prajwal (Single)	9	19.45
3	I <sub>3</sub>	Kodairoad (Dindugaldt)	Prajwal (Single)	14	34.33
4	I <sub>4</sub>	Kannanur (Madurai dt)	Prajwal (Single)	8	12.00
5	I <sub>5</sub>	Kannanur (Madurai dt)	Prajwal (Single)	12	24.65
6	I <sub>6</sub>	Sekanoorani (Madurai dt)	Prajwal (Single)	10	17.11
7	I <sub>7</sub>	Chellampatti (Madurai dt)	Prajwal (Single)	28	25.33



1. Pinpricked flower  
2. Without pricking  
3. Control

Figure 2: Pathogenicity on detached flower bud

## REFERENCES

- Arx, J.A. (1987), Plant Pathogenic Fungi. Cramer, Berlin.
- Bhattacharjee, S.K., Mukherjee, T. and Yadav, L.P. (1994), Standardization of agro-techniques in tuberose (*Polianthes tuberosa* Linn.) *Indian perfumer*, **38 (4)**: 144-152.
- Biswas B, Naveenkumar P and Bhattacharjee S K (2002), In: *Tuberose, AICRP on floriculture, Technical Bulletin*. No. 21.
- Burhan, M.J. (1987), *Botryodiplodia* on mango and banana. *FAO Pl. Prot. Bull.*, **35**: 34.
- Durgadevi D and Sankaralingam A (2012), First report of peduncle blight of tuberose caused by *Lasiodiplodiatheobromae* in India. *New disease reports*, **26**: 5.
- Khanzada, M. A, Rajput A. Q and Shahzad, S. (2006), Effect of medium, temperature, light and inorganic fertilizers on *in vitro* growth and sporulation of *Lasiodiplodiatheobromae* isolated from mango. *Pak. J. Bot.*, **38**: 885-889.
- Khanzada, M.A., Lodhi, A.M. and Shahzad, S. (2004), Mango die-back and gummosis in sindh, Pakistan caused by *Lasiodiplodiatheobromae*. *Plant Health Progress.*, **10**: 1094.
- Olunloyo, O.A. (1979), Efficiency of a single spray application of fungicide/ insecticide combination in the control of inflorescence blight disease of cashew. *Ann. Rept. Cocoa Res. Inst. Nigr.* p. 64-65.
- Patil, L.V., Shinde, V.B., Ghawade, R.S.D. and Wavare, S.H. (2006), Physiological and nutritional studies of *Botryodiplodiatheobromae* Pat.causing die-back disease of mango. *J. Pl. Dis. Sci.*, **1**: 216-218.
- Punithalingam, E. (1979), Plant Diseases attributed to *Botryodiplodiatheobromae*. *J. Cramer, Germany*. pp. 123.
- Wood, P.M. and Wood, C.E. (2005), Cane dieback of dawn seedless table grapevines (*Vitis vinifera*) in western Australia caused by *Botryosphaeria rhodina*. *Australas. Plant Pathol.*, **34**: 393-395.

