

ON THE SPECTRUM OF SELF-ADJOINT LÉVY GENERATORS

DAVID APPLEBAUM*

ABSTRACT. We investigate the spectrum of the generator of a self-adjoint transition semigroup of a (symmetric) Lévy process taking values in d -dimensional space.

1. Introduction

Lévy processes are essentially stochastic processes with independent increments. They have been extensively studied in recent years, as they include not only the most important Gaussian process – Brownian motion, but also many rich processes with jumps such as the α -stable ones. Four monograph treatments have been published in the last twenty five years that explore different aspects of their theory and application [1, 3, 12, 13]. The transition semigroups induced by the process may be studied in various Banach spaces, such as the space of continuous functions that vanish at infinity and the L^p -spaces. The fact that these operators, and their generators, may be written as pseudo-differential operators, whose symbol is given by the characteristic exponent of the process, has led to extensive work on Lévy-type operators that generate a very large class of Feller processes – see e.g. [4, 8, 9, 10, 11]. In this short note, we are interested in the generator as an operator acting in $L^2(\mathbb{R}^d)$.

For a long time, necessary and sufficient conditions have been known for when such an operator is self-adjoint and this material has recently been reviewed in Chapter 4 of [2]. Such operators are of interest in mathematical physics as they include both the free non-relativistic Hamiltonian (i.e. the (negative) Laplacian), and its relativistic counterpart. Some deep perturbation theory studies of these operators (at the form sum level), have been carried out in [7], while [5] has studied the relationship between recurrence of the underlying Lévy process and the existence of at least one negative bound state under perturbation by negative bounded potentials with compact support. However the author is not aware of any studies of the spectrum of the operator in question. For the case of Lévy processes on the torus, the spectrum consists of eigenvalues corresponding to the range of the characteristic exponent (see section 5.2.3 in [2]). When we work on \mathbb{R}^d , we expect to obtain a continuous spectrum which is again given by the closure of the range of the characteristic exponent. After collecting together all the known

Received 2019-5-30; Accepted 2019-6-13; Communicated by the editors.

2010 *Mathematics Subject Classification*. Primary 60G51; Secondary 47A10, 47G30.

Key words and phrases. Lévy generator, convolution semigroup, Lévy-Khintchine formula, Lévy measure, spectrum.

* Corresponding author.

facts we need in section 2, we will prove this conjecture in section 3 and present some examples. We use only elementary techniques in this note, the main purpose of which is to stimulate interest in the somewhat neglected area where spectral theory meets probability.

2. Preliminaries

The material in the section can be found in Chapters 3 and 4 of [2]. Much of it is also in Chapter 3 of [1]. Let $(\mu_t, t \geq 0)$ be a weakly continuous convolution semigroup of probability measures in \mathbb{R}^d . These arise naturally as the collections of laws of a Lévy process taking values in \mathbb{R}^d . We obtain a strongly continuous semigroup $(T_t, t \geq 0)$ of contractions in $L^2(\mathbb{R}^d)$ by the prescription

$$T_t f(x) = \int_{\mathbb{R}^d} f(x+y) \mu_t(dy),$$

for all $x \in \mathbb{R}^d, t \geq 0, f \in L^2(\mathbb{R}^d)$. In this article, we will require that μ_t is symmetric for all $t \geq 0$, i.e. $\mu_t(B) = \mu_t(-B)$, for all Borel sets B in \mathbb{R}^d . In this case the semigroup $(T_t, t \geq 0)$ is self-adjoint, and its self-adjoint infinitesimal generator A acts on the space $C_c^2(\mathbb{R}^d)$ of twice continuously differentiable functions with compact support on \mathbb{R}^d by the prescription

$$\begin{aligned} Af(x) &= \sum_{i,j=1}^d a_{ij} \partial_i \partial_j f(x) \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^d} (f(x+y) - 2f(x) + f(x-y)) \nu(dy), \end{aligned} \quad (2.1)$$

for all $f \in C_c^2(\mathbb{R}^d), x \in \mathbb{R}^d$, where $a = (a_{ij})$ is a non-negative definite symmetric $d \times d$ matrix, and ν is a symmetric Lévy measure, i.e.

$$\nu(\{0\}) = 0 \text{ and } \int_{\mathbb{R}^d} (|y|^2 \wedge 1) \nu(dy) < \infty,$$

(see Chapter 4 of [2] for the proof).

We have the Lévy–Khinchine formula

$$\int_{\mathbb{R}^d} e^{iu \cdot x} \mu_t(dx) = e^{-t\eta(u)},$$

where η is a continuous, negative definite function with $\eta(0) = 0$ that takes the form

$$\eta(u) = au \cdot u + \int_{\mathbb{R}^d} (1 - \cos(u \cdot y)) \nu(dy), \quad (2.2)$$

for all $u \in \mathbb{R}^d$. We define the usual Fourier transform \widehat{f} of $f \in L^2(\mathbb{R}^d)$ by

$$\widehat{f}(y) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} e^{-ix \cdot y} f(x) dx,$$

for $y \in \mathbb{R}^d$. The domain of the generator A is the non-isotropic Sobolev space

$$\mathcal{H}_\eta(\mathbb{R}^d) = \left\{ f \in L^2(\mathbb{R}^d), \int_{\mathbb{R}^d} \eta(y)^2 |\widehat{f}(y)|^2 dy < \infty \right\}.$$

Clearly $C_c^2(\mathbb{R}^d) \subseteq \mathcal{H}_\eta(\mathbb{R}^d)$, and on this larger domain we have the pseudo-differential operator representation

$$Af(x) = -\frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} e^{ix \cdot y} \eta(y) \widehat{f}(y) dy. \quad (2.3)$$

Underlying (2.3) is the formal relation

$$Ae^{iu \cdot x} = -\eta(u)e^{iu \cdot x}, \quad (2.4)$$

which suggests that the spectrum $\sigma(A) = \overline{\text{Ran}(-\eta)}$. This is certainly true when A is the Laplacian ($a_{ij} = \delta_{ij}$ and $\nu = 0$). We investigate it next in the more general context.

3. Results

Let B be a closed linear operator in a real Banach space E with domain $D_B \subseteq E$. We recall that its spectrum $\sigma(B)$ is the closed subset of \mathbb{C} defined by

$$\sigma(B) := \{\lambda \in \mathbb{C}; \lambda I - B \text{ fails to be invertible from } D_B \text{ to } E\}.$$

The resolvent set of B is $\rho(B) := \mathbb{C} \setminus \sigma(B)$. We need a general result. It is surely well-known, but we include a proof for the reader's convenience.

Proposition 3.1. *If $(S_t, t \geq 0)$ is a strongly continuous self-adjoint contraction semigroup acting in a Hilbert space H and having infinitesimal generator B then*

$$\sigma(B) \subseteq (-\infty, 0].$$

Proof. Since B generates a contraction semigroup, by the Hille–Yosida theorem, $\sigma(B) \subseteq \{z \in \mathbb{C}; \Re(z) \leq 0\}$, but B is self-adjoint and so $\sigma(B) \subseteq \mathbb{R}$. The result follows. \square

Our approach to investigating the spectrum of A is by a standard argument using Fourier transforms, as is well-known for the case of the Laplacian.

Theorem 3.2.

$$\sigma(A) = \overline{\text{Ran}(-\eta)}.$$

Proof. Consider the equation

$$(\lambda I - A)f = g,$$

for $\lambda \in \mathbb{C}, f \in \mathcal{H}_\eta(\mathbb{R}^d), g \in L^2(\mathbb{R}^d)$. Taking Fourier transforms of both sides we obtain

$$(\lambda + \eta(y))\widehat{f}(y) = \widehat{g}(y),$$

for all $y \in \mathbb{R}^d$. If the operator equation has a solution, then

$$(1 + \eta(\cdot))\widehat{f}(\cdot) = \frac{1 + \eta(\cdot)}{\lambda + \eta(\cdot)}\widehat{g}(\cdot) \in L^2(\mathbb{R}^d).$$

This clearly cannot be the case when $\lambda = -\eta(y)$ for some $y \in \mathbb{R}^d$, and it follows that $\overline{\text{Ran}(-\eta)} \subseteq \sigma(A)$.

Our goal now is to prove that the resolvent set $\rho(A) = \mathbb{C} \setminus \overline{\text{Ran}(-\eta)}$. By Proposition 3.1, we know that $\mathbb{C} \setminus (-\infty, 0] \subseteq \rho(A)$. So it is sufficient to consider

$\lambda \in (-\infty, 0) \setminus \text{Ran}(-\eta)$. The required result holds if g_λ is bounded, where for all $u \in \mathbb{R}^d$,

$$g_\lambda(u) := \frac{1 + \eta(\cdot)}{\lambda + \eta(\cdot)}.$$

There are two cases to consider. Firstly assume that $\sup_{u \in \mathbb{R}^d} \eta(u) = K < \infty$. Then $\text{Ran}(\eta) = [0, K]$. Choose $\epsilon > 0$ and assume that $\lambda \in (\infty, -K - \epsilon)$. Then we find that

$$\sup_{u \in \mathbb{R}^d} |g_\lambda(u)| \leq \frac{1 + K}{\epsilon} < \infty.$$

But then we conclude that

$$(-\infty, -K) = \bigcup_{\epsilon > 0} (-\infty, -K - \epsilon) \subseteq \rho(A),$$

and so $\sigma(A) = \overline{\text{Ran}(-\eta)}$, as required. If $\sup_{u \in \mathbb{R}^d} \eta(u) = \infty$, then by the first part of the proof we have $(-\infty, 0) \subseteq \sigma(A)$ and so the required result follows directly from Proposition 3.1. \square

Corollary 3.3. *If $\lim_{u \rightarrow \infty} \eta(u) = \infty$, then $\sigma(A) = (-\infty, 0]$.*

Proof. This is an immediate consequence of Theorem 3.2 and the continuity of η . \square

We note that the condition of Corollary 3.3 was also required in [5] to obtain the result alluded to in the introduction.

Some well-known and important examples satisfy the conditions of Corollary 3.3. These include

- The Laplacian Δ , where $\eta(u) = |u|^2$.
- The fractional Laplacian $-(-\Delta)^{\alpha/2}$, where $\eta(u) = |u|^\alpha$ with $0 < \alpha < 2$.
- The relativistic Schrödinger operator $-\sqrt{b^2 I - \Delta} + bI$, where $\eta(u) = \sqrt{b^2 + |u|^2} - b$ for some $b > 0$.

More generally, to obtain $\sigma(A) = (-\infty, 0]$ we may take $A = -f(-\Delta)$, where f is a strictly increasing Bernstein function for which $f(0) = 0$, in which case the underlying Lévy process is a subordinated Brownian motion (see e.g. [14]). The above three examples are all special cases of this class. In the case where f is a complete Bernstein function, our main result Theorem 3.2, is a special case of Theorem 13.11 in [14] (p.211).

To find other interesting class of examples, we first write $A = A_1 + A_2$, where A_1 is the second order differential operator in (2.1).

Lemma 3.4. *If ν is a finite measure and $M := \nu(\mathbb{R}^d)$, then A_2 is a bounded operator in $L^2(\mathbb{R}^d)$, with $\sigma(A_2) \subseteq [-2M, 0]$.*

Proof. Since the measure ν is finite and symmetric, $A_2 f(x) = \int_{\mathbb{R}^d} (f(x+y) - f(x)) \nu(dy)$, for $f \in L^2(\mathbb{R}^d)$, $x \in \mathbb{R}^d$. For each $y \in \mathbb{R}^d$, define the shift operator τ_y by $\tau_y f(x) = f(x+y)$. The linear operator A_2 is bounded as by convexity of the

norm, Jensen's inequality and unitarity of τ_y , we have

$$\begin{aligned} \|A_2 f\| &\leq \int_{\mathbb{R}^d} \|\tau_y f + f\| \nu(dy) \\ &\leq 2M \|f\|, \end{aligned}$$

The second result follows from the fact that for all $u \in \mathbb{R}^d$

$$\eta(u) = \int_{\mathbb{R}^d} (1 - \cos(u \cdot y)) \nu(dy) \leq 2M.$$

□

A necessary and sufficient condition for $\sigma(A_2) = [-2M, 0]$ is that

$$\nu(\{y \in \mathbb{R}^d; u \cdot y = (2n+1)\pi \text{ for some } u \in \mathbb{R}^d, n \in \mathbb{Z}\}) > 0.$$

Now let $\alpha > 0$ be arbitrary. For an example where $\sigma(A_2) = [-\alpha, 0]$, take $d = 1$ and $\nu = \frac{\alpha}{4}(\delta_{-1} + \delta_1)$.

Corollary 3.5. *If ν is finite and there exist $b_i > 0$ for all $i = 1, \dots, d$ such that*

$$au \cdot u \geq \sum_{i=1}^d b_i u_i^2$$

for all $u \in \mathbb{R}^d$, then $\sigma(A) = (-\infty, 0)$.

Proof. This follows from the fact that $\eta(u) \geq \sum_{i=1}^d b_i u_i^2 \rightarrow \infty$, as $u \rightarrow \infty$. □

Note that the condition in the last corollary is satisfied if a is bounded below in the sense that there exists $b > 0$ so that $a \geq bI$.

Remark 3.6. As expected, both the semigroup operators and generator are unitarily equivalent to multiplication operators, and the unitary equivalence is implemented by the Fourier transformation $\mathcal{F} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$, where $\mathcal{F}(f) = \widehat{f}$ for each $f \in L^2(\mathbb{R}^d)$. Then if we define $\eta(D)\widehat{f}(u) = \eta(u)\widehat{f}(u)$ for all $f \in \mathcal{H}_\eta(\mathbb{R}^d)$, $u \in \mathbb{R}^d$, we have

$$Af = \mathcal{F}^{-1}(-\eta(D))\mathcal{F}f,$$

and (with an obvious notation), for all $t \geq 0$.

$$\mathcal{F}T_t\mathcal{F}^{-1} = e^{-\eta(D)t}.$$

This gives rigorous meaning to the formal relation (2.4).

Acknowledgment. I am grateful to René Schilling for helpful comments on the first draft of this paper, and the referee for correcting some typos.

References

1. Applebaum, D.: *Lévy Processes and Stochastic Calculus* (second edition), Cambridge University Press (2009)
2. Applebaum, D.: *Semigroups of Linear Operators with Applications to Analysis, Probability and Physics*, to appear Cambridge University Press (2019)
3. Bertoin, J.: *Lévy Processes*, Cambridge University Press (1998)
4. Böttcher, B., Schilling, R. L., and Wang, J.: *Lévy Matters III, Lévy Type Processes, Construction, Approximation and Sample Path Properties*, Lecture Notes in Mathematics Vol. **2099**, Springer International Publishing, Switzerland (2013)
5. Carmona, R., Masters, W. C., and Simon, B.: Relativistic Schrödinger operators; asymptotic behaviour of the eigenfunctions, *J. Funct. Anal.* **91** (1990), 117–42
6. Engel, K.-J., Nagel, R.: *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag New York Inc. (2000)
7. Herbst, I. R. and Sloan, A. D.: Perturbation of translation invariant positivity preserving semigroups on $L^2(\mathbb{R}^n)$, *Trans. Amer. Math. Soc.* **236** (1978), 325–60
8. Jacob, N.: *Pseudo-Differential Operators and Markov Processes*, Akademie-Verlag, Mathematical Research 94 (1996)
9. Jacob, N.: *Pseudo-Differential Operators and Markov Processes: 1, Fourier Analysis and Semigroups*, World Scientific (2001)
10. Jacob, N.: *Pseudo-Differential Operators and Markov Processes: 2, Generators and Their Potential Theory*, World Scientific (2002)
11. Jacob, N.: *Pseudo-Differential Operators and Markov Processes: 3, Markov Processes and Applications*, World Scientific (2005)
12. Kyprianou, A. E.: *Fluctuations of Lévy Processes with Applications*, Springer-Verlag Berlin Heidelberg (2014)
13. Sato, K.-I.: *Lévy Processes and Infinite Divisibility*, Cambridge University Press (1999)
14. Schilling, R. L., Song, R., and Vondraček, Z.: *Bernstein Functions, Theory and Applications*, Studies in Mathematics **37**, De Gruyter (2010), second edition (2012)
15. Mansor H. Ibrahim., International Financial Integration and Real Economic Activity in A Small Open Economy, *The Global Journal of Finance and Economics*, Vol. 3, No. 1, pp. 17-37.
16. Mukhriz Izraf Azman Aziz & Nor’Aznin Abu Bakar, Oil Price Shocks and Macroeconomic Activities in Malaysia, *Journal of World Economic Review*
17. Adam Bouras, David Bouras and Emmanuel Ajuzie, An Empirical Analysis of the effect of Diversification on Financial Performance: The Case of the US Grocery Store Industry, *Review of Applied Economics*
18. Lefteris Tsoulfidis & Theologos Dergiades., Time Series Estimates of Capacity Utilization: A Profits and Investment based Approach, *International Journal of Economics*, Vol. 1, No. 1, pp. 63-74.
19. Ren-huai Liu., Large Deflection of Annular Sandwich Plates, *Journal of Mechanics and MEMS*
20. Bellanca Sabrina & Cultrera Loredana, The Accounting Reform in the Brussels-Capital Region: New Tool for Financial Analysis, *Global Review of Business and Economic Research*
21. Kamal Sarkar: Application of Dimensional Analysis to Predict Poly Ethylene Oxide (PEO) Fiber Diameters from Electrospinning Process, *International Journal of Electrospun Nanofibers and Applications*
22. Fregly Benjamin : Design of Optimal Treatments for Neuromusculoskeletal Disorders using Patient-Specific Multibody Dynamic Models, *International Journal of Computational Vision and Biomechanics*, Vol. 2 No. 2 (2016)
23. Everton Dockery., Asymmetries and the Effect of Macroeconomic News on Stock Market Volatility: An Empirical Examination of the Warsaw Equity Market, *International Journal of Financial Economics and Econometrics*, Vol. 1, No. 1, pp 11-29.
24. Lan LUO, Notes to Grain128_p2 of ESTREAM Phases 3, *International Journal of Computer Science, Information Technology and Management*
25. Mahender Singh, P. C. Basak & Rajbir Singh., Competitive Factors and Their Relative Importance in the Indian Manufacturing Industries, *Journal of Asian Business Management*, Vol. 1, No. 1, pp. 13-18.
26. Mohammad Hossein Alipour , A NEW FUZZY TECHNIQUE TO FIND THE OPTIMAL SOLUTION IN FLOOD MANAGEMENT, *Journal of Flood Engineering*
27. Tatjana Volkova, VALUE INNOVATION FRAMEWORKS FOR DELIVERING SUPERIOR CUSTOMER RESPONSIVENESS, *Journal of Business Management*
28. Yanil Zhang, Resource Based and Knowledge Based Views as Complementary Perspectives to Transaction Cost Theory, *International Journal Data Modelling and Knowledge Management*
29. D. Paschaloudis, K. Anastasiadou, A. Anastasiadou & P. Pantelidis., The Internationalisation Paths,

- International Review of Applied Economic Research*, Vol. 1, No. 1, pp. 23-34.
30. Martin Burda, Roman Liesenfeld, and Jean-Francois Richard, Bayesian Analysis of a Probit Panel Data Model with Unobserved Heterogeneity and Autocorrelated Errors, *International Journal of Statistics and Management System*
 31. Xia Mu, A Hybrid Approach for Mobile Agent Security using Reversible Watermarking, *Journal of Information Technology and Engineering*
 32. Al-Lawatia, Numerical Studies for One-dimensional Nonlinear Coupled System of Parabolic and Hyperbolic Equations, *International Journal of Computational Mathematics and Numerical Simulation*
 33. A. Jafaripour, Analytical Solution of the Falkner–Skan Equation for Wedge, *International Journal of Nonlinear Dynamics in Engineering and Sciences*
 34. Tain-Sou Tsay, On-Line Computing and Control for Decoupling Multivariable Processes with Gain and Phase Specifications, *International Journal of Contemporary Mathematics*

D.APPLEBAUM: SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SHEFFIELD, HICKS BUILDING, HOUNSFIELD ROAD, SHEFFIELD S3 7RH, ENGLAND
E-mail address: D.Applebaum@sheffield.ac.uk